

JBUMS

# Comparison of the Dental Caries Experience and Some Salivary Physicochemical Variables among Iraqi Boxers

H. Obied Majhool (DDS, MS)<sup>\*1</sup><sup>(b)</sup>, A. Ahmed Qasim (DDS, MS)<sup>1</sup><sup>(b)</sup>

1. Department of Preventive Dentistry, College of Dentistry, University of Baghdad, Baghdad, Iraq.

| *Corresponding Author     | or: H. Obied Majhool (DDS, MS)  |
|---------------------------|---|
| Address: Department of    | of Preventive Dentistry, College of Dentistry, University of Baghdad, Baghdad, Iraq.                          |
| Tel: +964 (783) 05233     | 37. E-mail: haneen93obied@gmail.com   |
| Article Type              | ABSTRACT  |
| <b>Research Paper</b>     | Background and Objective: Due to the importance of oral hygiene to athletes in the evaluation of              |
|                           | oral health status, efforts continue to raise their performance. This study aims to determine a               |
|                           | correlation between some salivary physicochemical variables to dental caries in Iraqi boxers.                 |
|                           | Methods: This cross-sectional study was conducted on 80 people in two groups of 40, including 40              |
|                           | boxers in different boxing clubs in Baghdad and 40 male students aged 15-16 from different schools            |
|                           | in Baghdad. The two groups were compared in terms of some variables including age, sex, education             |
|                           | status, general health status. The clinical assessment of dental caries was done by dental mirror and         |
|                           | dental probes. The unstimulated salivary analysis included samples collected to determine pH                  |
|                           | (measured by pH meter), flow rate (By dividing the measured amount of saliva by ml/min) and                   |
|                           | minerals like calcium (measured by calcium kit), and salivary immunoglobulin A (measured by                   |
|                           | ELISA kit).   |
|                           | Findings: There were no significant differences between the two groups in terms of demographic                |
|                           | variables. Dental caries was seen in the study group in 100% of 40 subjects compared to control               |
|                           | group (2.5%). The mean value of decay surfaces, missing surfaces, decay missing filling teeth, and            |
|                           | decay missing filling surfaces show high levels with a significant difference in the study group              |
| Received:                 | 20.275±6.880, 4.050±5.514, 11.775±3.779 and 24.450±9.375 respectively (p<0.05). The mean                      |
| Jun 11 <sup>st</sup> 2023 | values to the pH and flow rate were $5.903\pm0.568$ , $0.939\pm0.445$ respectively (p<0.05). For the salivary |
| Davisod.                  | immunoglobulin A (IgA) and calcium, they were about 0.375±0.123, 0.887±0.260 respectively                     |
| Neviseu.                  | (p<0.05) and were lower in the study groups, indicating statistically significant differences.                |
| Jul 10 <sup>th</sup> 2023 | Conclusion: According to the result of this study, dental caries had a negative correlation with pH           |
| Accepted:                 | and flow rate. Also, salivary calcium and salivary IgA show a negative correlation with dental caries.        |
| Jul 26 <sup>th</sup> 2023 | Keywords: Iraqi Boxer, Dental Caries, Salivary Physicochemical Variables.                                     |
| Cita this article: Ob     | ied Maihool H. Ahmed Asim A. Comparison of the Dental Caries Experience and Some Salivary                     |

**Cite this article:** Obied Majhool H, Ahmed Qasim A. Comparison of the Dental Caries Experience and Some Salivary Physicochemical Variables among Iraqi Boxers. *Journal of Babol University of Medical Sciences*. 2024; 26: e63.



## Introduction

The most common kind of chronic oral disease that mostly harms the health of humans is dental caries (1). Despite the progress in oral health in recent years, there stays oral health problems globally due to it being widely infectious and widespread (2, 3). Oral diseases are highly significant and the WHO analysis reported that they affect 621 million teenagers globally. In teenagers, the general standard decayed-missingfilled teeth (DMFT) is about 1.86.3. The communication of a mass of cariogenic oral microorganisms can lead to caries (4, 5). A high risk of dental caries is reported in athletes in different studies; about 78.72 % (6) and 49.1% (7). According to the findings, good nutritional health leads to good oral health (8). Sports dentistry seeks to reply to these athletes' particular requirements (9). Saliva can determine oral health status by salivary components (10). Many researchers used salivary diagnostics as a precious tool for systemic conditions in diagnosis (11). Saliva flow rate is the volume of Saliva which changes during lifespan (12). Stress may be connected with several pathologies of the oral cavity (13). Oral and dental health including caries may result from the negative impact of stressful life (14). A significant positive association is seen between salivary pH and flow rate (15). Many of the electrolytes in the body diminish. As a result, sports performance is negatively influenced by insufficient electrolytes. Sports scientists were interested in it due to its impact on athletic performance (16). The large appropriate immune constituent produced by mature B lymphocytes is Secretory IgA (17).

Therefore, the objective of this research is to detect the exercise-related factors like stress, dry mouth, and fatigue that may lead to changes in oral diseases, including dental caries and salivary variables among boxers and to evaluate the correlation between Iraqi boxers' dental caries and some physicochemical salivary variables. More studies are required to give them the care specialty in oral health they meet and to dissolve all their problems.

## Methods

After approval by the ethics committee of the University of Baghdad's College of Dentistry on 2/6/2022, Ref. number: 579, this cross-sectional comparative was conducted among 80 males, including 40 male boxers with weights ranging from (46-60 kg) and 40 male school students, in the age range of (15-16) as case and control groups, respectively. The study was conducted in different boxing clubs in Baghdad and different schools in Baghdad. The sample size of this study was measured by utilizing G power 3.1.9.7 (Program by Franz- Faul, Kiel University, Germany), alpha error of probability=0.05, study power=85%, Independent T-test and assuming the effect size of 0.6 in the two groups. So, the sample size was 40 for both groups. Cohen's d was as follows: small=0.3, medium=0.5, large $\geq$ 0.8. This number includes only Baghdad in Iraq. Clinical and demographic data were also collected. These include sex, age, education status and practicing in boxing clubs. We exclude people with systemic disease, uncooperative people, smokers, hormonal stimulations, and if they did other sports.

The procedure was done by oral examination and was carried out according to the basic method of an oral health survey by the WHO 2013. The four surfaces of each tooth were examined. All teeth were included except the third molars (18). Clinical examinations were done using dental mirrors and dental probes. To collect unstimulated salivary samples, the people were ordered to avoid food, smoke, and beverages 1 hour before saliva collection. The subject then washed his mouth with distilled and rested for five minutes. A subject had to fix the forehead over the test tube and keep steady. To collect Saliva in the tube for five minutes, subjects were required to keep their mouth open. In the last stage, the subject was told to keep any remaining saliva in the mouth and immediately spit it in the exam tube. The actual timespan

must be almost 5 minutes. Then, when saliva collection was completed and the foam disappeared, the flow rate of saliva was detected by dividing all collections of saliva in milliliters by the minutes consumed until the sample was fixed (10 min), expressed by ml/min (11). Flow rate= Volume (ml)/Time (min). Saliva pH was measured by merging the pH meter's sensor into the tube containing saliva and the tube was tilted so that the saliva could cover the entire surface of the sensor. Saliva was sent to the lab in a cooling box. Then, samples were centrifuged at 3000 rpm for 10 minutes. Then a micropipette was used to separate the supernatant and underwent deep freezing (-20°C). The Salivary calcium concentration released in the demineralizing solution was measured by spectrophotometry in a lab. Then the standards, samples, and quality control specimens were diluted according to instructions 50-fold with lanthanum diluents, mix, and calcium measured by using AAS. A hollow cathode lamp at a wavelength of 422.7 nm specific was used for calcium. ELISA kit was used for determent concentration of immunoglobulin IgA level (enzyme-linked immunosorbent assay). The required streaks dictated by the samples were as follows:

1-Put in only Chromogen solutions A with B, and break off fluid.

2-Put in 50 µl standard and streptavidin-HRP 50µl.

3-Put in 40 µl of sample and then 10µl IgA antibodies, 50 µl streptavidin-HRP to sample well to be tested.

Then with a seal plate membrane, they were strongly shaken to mix them up. Incubation was done at 37°C for 60 minutes. Fluid prepared to wash: it was debilitated with distilled water to washing concentration (30X) for later use. After washing, seal paten membrane was used, the solution was drawn off, and the remaining liquid was vibrated, and all wells were cleaned with a wiping solution. The liquid drawn off after 30 seconds of staying still. This process was repeated 5 times. Evaluation color: Put in 50µl chromogen solution A to all well and then put in 50µl liquid of chromogen B to all. Vibrate firmly to mix them up. Incubate at 37°C for 10 minutes away from light for color Evaluation. Then put in 50µl liquid over both well to be over that short time the reaction changed from color blue to yellow.

Data analysis, presentation, and description were conducted by Statistic Package to Social Sciences SPSS version 22, Chicago, Illinois USA, Statistic analyses categorized into descriptive analysis consist of frequency, and percentage for qualitative variables while mean, standard error (SE) and standard deviation (SD) for quantitative variable. Graphs included cluster chart bars and the analysis inferential as independent sample T-test parametric test the difference between two groups, human correlation: parametric test for correlation between two quantitative variables and Levene test for homogeneity of variance in groups.

#### Results

The study shows that dental caries was high in study group (100% of the 40 people). The mean value of decay surfaces, missing surfaces, decay missing filling teeth, and decay missing filling surfaces was higher in study group with a significant difference of  $20.275\pm6.880$ ,  $4.050\pm5.514$ ,  $11.775\pm3.779$  and  $24.450\pm9.375$ , respectively (p<0.05). Except for FS; it was significantly different between the two groups, and lower in the study group as seen in Table 1.

The study group showed increase in salivary flow rate and pH with significant differences (p<0.05), and the mean values were  $0.939\pm0.445$ ,  $5.903\pm0.568$ , respectively (Table 2). The mean values of salivary immunoglobulin IgA and calcium in the study group were  $0.375\pm0.123$ ,  $0.887\pm0.260$ , respectively (p<0.05), and found to be low with significant difference (Table 3).

The result of this relation includes that there was significant negative correlation between PH and salivary flow rate and dental caries in the study group (-0.540, -0.134, -0.500, -0.170), respectively (p<0.05). At the same time, other results are weak, with no significant correlations, as seen in Table 4.

There was a weak significant negative correlation between salivary IgA and calcium with the missing surface in the control group (-0.245, -0.407) respectively (p<0.05), while other results are not significant (Table 5).

| Table 1. Descriptive of caries experience |                    |       |                     |       |        |         |  |
|---|--------------------|-------|---------------------|-------|--------|---------|--|
|   |                    | Gr    | oups                |       |        |         |  |
| Vars.                                     | Study              |       | Control             |       | T-test | p-value |  |
|   | <b>Mean±SD</b>     | ±SE   | Mean±SD             | ±SE   |        |         |  |
| DS  | $20.275 \pm 6.880$ | 1.088 | 15.675±10.247       | 1.620 | 2.357  | 0.021   |  |
| MS  | $4.050 \pm 5.514$  | 0.872 | $1.000 \pm 2.025$   | 0.320 | 3.284  | 0.002   |  |
| FS  | $0.125 \pm 0.463$  | 0.073 | $0.975 \pm 1.423$   | 0.225 | 3.592  | 0.001   |  |
| DMFT                                      | 11.775±3.779       | 0.598 | $9.550 \pm 5.602$   | 0.886 | 2.082  | 0.041   |  |
| DMFS                                      | $24.450 \pm 9.375$ | 1.482 | $17.650 \pm 10.888$ | 1.721 | 2.993  | 0.004   |  |

Table 2. Physicals constituent salivary pH with a flow rate

| Vars. and groups | Mean±SD           | ±SE   | T-test  | p-value  |
|------------------|-------------------|-------|---------|----------|
| рН               |                   |       |         |          |
| Study            | 5.903±0.568       | 0.090 | 2 1 4 4 | 0.025    |
| Control          | 6.225±0.763       | 0.121 | 2.144   | 0.055    |
| FR               |                   |       |         |          |
| Study            | 0.939±0.445       | 0.070 | 2 050   | 0.000224 |
| Control          | $1.248 \pm 0.241$ | 0.038 | 3.838   | 0.000234 |

Table 3. Chemical constituents salivary IgA and calcium Groups Vars. Study T-test p-value Control **Mean±SD** ±SE Mean±SD ±SE SIGA  $0.375{\pm}0.123 \quad 0.019 \quad 0.420{\pm}0.035 \quad 0.006$ 2.222 0.031 Calcium 0.887±0.260 0.041 0.989±0.139 0.022 2.181 0.033

| Table 4. Correlation between caries, pH, and salivary flow |        |       |        |       |       |       |        |       |        |       |
|--|--------|-------|--------|-------|-------|-------|--------|-------|--------|-------|
| Crong  | D      | S     | MS     |       | FS    |       | DMFT   |       | DMFS   |       |
| Groups   | R      | Р     | R      | Р     | R     | Р     | R      | Р     | R      | Р     |
| Control  |        |       |        |       |       |       |        |       |        |       |
| pН   | -0.124 | 0.445 | -0.008 | 0.960 | 0.038 | 0.817 | -0.141 | 0.385 | 0.096  | 0.557 |
| FR   | -0.245 | 0.127 | -0.204 | 0.207 | 0.021 | 0.900 | -0.210 | 0.194 | -0.285 | 0.075 |
| Study  |        |       |        |       |       |       |        |       |        |       |
| pН   | -0.524 | 0.001 | -0.187 | 0.249 | 0.140 | 0.389 | -0.540 | 0.000 | -0.500 | 0.001 |

| Crouns  | SIGA   |       | Phosp  | horous | Calcium |       |  |
|---------|--------|-------|--------|--------|---------|-------|--|
| Groups  | R      | Р     | R      | Р      | R       | Р     |  |
| Control |        |       |        |        |         |       |  |
| DS      | -0.386 | 0.014 | -0.218 | 0.177  | 0.028   | 0.864 |  |
| MS      | -0.245 | 0.128 | -0.023 | 0.886  | 0.407   | 0.009 |  |
| FS      | 0.011  | 0.947 | -0.114 | 0.484  | -0.236  | 0.143 |  |
| DMFT    | -0.352 | 0.026 | -0.070 | 0.669  | -0.159  | 0.327 |  |
| DMFS    | -0.410 | 0.009 | 0.082  | 0.615  | -0.133  | 0.413 |  |
| Study   |        |       |        |        |         |       |  |
| DS      | 0.018  | 0.913 | -0.141 | 0.385  | -0.043  | 0.794 |  |
| MS      | -0.230 | 0.154 | -0.073 | 0.653  | 0.072   | 0.661 |  |
| FS      | 0.077  | 0.635 | 0.038  | 0.814  | -0.034  | 0.835 |  |
| DMFT    | -0.153 | 0.347 | -0.075 | 0.647  | 0.024   | 0.883 |  |
| DMFS    | -0.118 | 0.468 | -0.062 | 0.703  | -0.072  | 0.660 |  |

Table 5. Correlation between salivary SIGA, calcium, and caries

#### Discussion

In this study, dental caries increased in study groups compared to control groups. This finding agrees with Whitaker et al. who reported the risk of dental caries is raised in athletes (19). This may come from the mouth guard outside the mouth, increasing the risk of contamination which leads to dental caries (20). This study showed low salivary pH and flow rate with significant differences in a study group. This may be related to the intensity of exercise as mouth breathing which leads to the salivary flow rate becoming low (21). Also, it may be from low flow rate in athletes, associated with effects of oral hygiene (22). This finding agrees with Mosca et al. who reported alteration in salivary pH concerning strong evidence of eating habits (23). The intensity of sport may cause changes in pH (22). They also showed that the salivary IgA and calcium in the study group were lower. This may be due to the low mineral levels in athletes which is connected with psycho-emotional and physical feelings (24). The salivary IgA is lower, which may agree with Rico-González et al. (25).

In study of Wu et al., the negative correlation between salivary flow rate and pH with caries (26), which coincides with the result of this study, explained the indicator of caries susceptibility to salivary flow rate. But some studies disagree, suggesting that dental caries do not affect salivary flow rate (27). Also, this may be explained by the time of decrease in salivary flow leading to dental caries (28). Regarding pH, this result agrees with the study of Wu et al. (26) who reported when dental caries increase, the pH decreases. It may also be related to the connection between physical training and carbohydrates, which leads to low salivary pH and causes dental caries (29). Also, the protective salivary constituent leads to a pH fall and develops dental caries (30). This study showed a negative correlation between calcium and dental caries. This may be connected with the study of Ram et al. (31) who reported that increase in dental caries is associated with decreased salivary calcium in exercise. This also explains that poeple with more dental caries have low ion in saliva (32). Also, this study shows a negative significant relation between IgA with dental caries. This agrees with study of Leary et al. (33) who reported the low IgA due to the athletes having poor oral health which increases dental caries.

It disagrees the with study of Gómez et al. (34) who illustrated low levels of salivary IgA with increased caries activity. Maybe the IgA can fight oral cavity microbes; therefore, when it decreases, it is associated with more caries (35). Also, maybe the IgA falls through training and causes increase in dental caries (36).

Based on the result of this study, high-intensity training, poor oral hygiene, and mouth breathing during exercise are connected with an increased prevalence of dental caries among boxers (100%). In the study group, PH and salivary flow rate showed significant differences. Salivary IgA and calcium are low in the study group, with significant differences. Considering this correlation, the result shows a negative correlation between dental caries and salivary flow rate among study players, and positive correlation between salivary IgA, and calcium in a study group with dental caries with significant differences.

## Acknowledgment

We would like to thank the Vice President for Research of Babol University of Medical Sciences and the Iraqi boxing federation and all people who participated in this study.

### References

1. Cheng L, Zhang L, Yue L, Ling J, Fan M, Yang D, et al. Expert consensus on dental caries management. Int J Oral Sci. 2022;14(1):17.

2. Alaei Bakhsh MA, Khosravi Samani M, Motalebnejad M, Bijani A, Ghorbani A, Mehryari M. Frequency of Oral Lesions and Its Relationship with Some Variables. J Babol Univ Med Sci. 2021;23:345-51. [In Persian]

3.Al-Obaidi ZQM, Radhi NJM. Does Gluten Free Diet and Delay in Celiac Disease Diagnosis Affect Dental Caries and Salivary Oxidative Stress in Children?. Indian J Public Health Res Dev. 2019;10(10):2098.

4.Martins MT, Sardenberg F, Bendo CB, Abreu MH, Vale MP, Paiva SM, et al. Dental caries remains as the main oral condition with the greatest impact on children's quality of life. PLoS One. 2017;12(10):e0185365.

5. Jaddoa MF, Qasim AA. Assessment of the Correlation between the Salivary Flow Rate and Dental Caries Experience among Children with  $\beta$ -Thalassemia Major. Indian J Forensic Med Toxicol. 2020;14(1):310-5.

6.Lima IAB, Quaresma FRP, Barasuol AM, Almeida RJL, Sousa LS, Pontes-Silva A, et al. Epidemiological study on dental caries in schoolchildren: A cross-sectional study. Res Soc Dev. 2023;12(3):e28212340871.

7.Gallagher J, Ashley P, Petrie A, Needleman I. Oral health and performance impacts in elite and professional athletes. Community Dent Oral Epidemiol. 2018;46(6):563-8.

8.AL-Nassary HTH, Mohammed AT. Dental Caries and Treatment Needs in Relation to Nutritional Status among Kindergartens Children in Tikrit City, Iraq. Indian J Public Health Res Dev. 2019;10(10):1742.

9.Sajjanar A, Rojekar N, Kumar S, Gahlod N, Shukla H, Bhattad D. Sports Dentistry: A Review. Ann Rom Soc Cell Biol. 2021;25(4):2031-7.

10.Awn BH, Munther S, Yas BA. Salivary Protein Carbonyl Level in Relation to Gingival Health Status among a Group of Iraqi Pregnant Women. Ann Trop Med Public Health. 2020;23(11):1129.

11.Huang Z, Yang X, Huang Y, Tang Z, Chen Y, Liu H, et al. Saliva - a new opportunity for fluid biopsy. Clin Chem Lab Med. 2022;61(1):4-32.

12.Xu F, Laguna L, Sarkar A. Aging-related changes in quantity and quality of saliva: Where do we stand in our understanding?. J Texture Stud. 2019;50(1):27-35.

13.Kadhum RI, Qasim AA. The impact of dental environment stress on caries experience, salivary flow rate and uric acid. J Bagh Coll Dent. 2020;32(1):35-41.

14. Abedi Gh, Esmaeili Seraji Z, Mahmoodi Gh, Jahani MA, Abbasi M. Evaluating the Implementation of Family Physician Program in Urban and Rural Areas of Mazandaran Province Based on Process Approach. J Babol Univ Med Sci. 2020;22(1):126-34. [In Persian]

15.Abdul-Ameer AK, Radhi NJ, Abdul-Ghani HJ. Stressful Life Events In Relation to Dental Caries and Selected Salivary Constituents among Secondary School Students in Baghdad City. J Bagh Coll Dent. 2017;29(1):139-47.

16.Jain S, Bansal K, Marwaha M, Sehrawat N, Singla S. Effect of Diet Modification on Salivary Parameters and Oratest in High-caries-risk Individuals. Int J Clin Pediatr Dent. 2018;11(1):34-9.

17.Brown EL, Essigmann HT, Hoffman KL, Palm NW, Gunter SM, Sederstrom JM, et al. Impact of Diabetes on the Gut and Salivary IgA Microbiomes. Infect Immun. 2020;88(12):e00301-20.

18. World Health Organization. Oral health surveys: basic methods, 5<sup>th</sup> ed. World Health Organization; 2013. Available from: https://iris.who.int/handle/10665/97035

19.Whitaker EJ, Al Anezi NS, Yesilsoy C. Evaluation of Hand Sanitizer for On-Site Disinfection of Mouthguards. Health. 2019;10(5):555796.

20.Duan Z, Qin J, Li C, Ying C. Clinical and Molecular Epidemiology of *Stenotrophomonas maltophilia* in Pediatric Patients From a Chinese Teaching Hospital. Front Cell Infect Microbiol. 2020;10:411.

21.Brancher JA, Morodome F, Madalena IR, Reis CLB, Von Held R, Antunes LAA, et al. Salivary pH and oral health of Brazilian para-athletes: Saliva and oral health of para-athletes. Spec Care Dentist. 2021;41(4):505-11.

22.McKay AKA, Pyne DB, Peeling P, Sharma AP, Ross MLR, Burke LM. The impact of chronic carbohydrate manipulation on mucosal immunity in elite endurance athletes. J Sports Sci. 2019;37(5):553-9.

23.Mosca AC, Stieger M, Neyraud E, Brignot H, van de Wiel A, Chen J. How are macronutrient intake, BMI, ethnicity, age, and gender related to the composition of unstimulated saliva? A case study. J Texture Stud. 2019;50(1):53-61.

24.Vdovenko NV, Ivanova AM, Loshkarova IA. Practical recommendations concerning prevention and correction of iron deficit in athletes. Pedagogics Psychol Med-Biol Probl Phys Train Sport. 2015;19(1):12-6.

25.Rico-González M, Clemente FM, Oliveira R, Bustamante-Hernández N, Pino-Ortega J. Part I: Relationship among Training Load Management, Salivary Immunoglobulin A, and Upper Respiratory Tract Infection in Team Sport: A Systematic Review. Healthcare (Basel). 2021;9(4):366.

26.Wu Z, Gong Y, Wang C, Lin J, Zhao J. Association between salivary s-IgA concentration and dental caries: A systematic review and meta-analysis. Biosci Rep. 2020;40(12):BSR20203208.

27.Dewi SRP, Septhimoranie S, Muchzal S. Correlation of saliva characteristics and caries in beta-thalassemia major patients. Majalah Kedokteran Gigi Indonesia. 2020;6(2):100-5.

28.Broughton D, Fairchild RM, Morgan MZ. A survey of sports drinks consumption among adolescents. Br Dent J. 2016;220(12):639-43.

29.Frese C, Frese F, Kuhlmann S, Saure D, Reljic D, Staehle HJ, et al. Effect of endurance training on dental erosion, caries, and saliva. Scand J Med Sci Sports. 2015;25(3):e319-26.

30.Pyati SA, Naveen Kumar R, Kumar V, Praveen Kumar NH, Parveen Reddy KM. Salivary Flow Rate, pH, Buffering Capacity, Total Protein, Oxidative Stress and Antioxidant Capacity in Children with and without Dental Caries. J Clin Pediatr Dent. 2018;42(6):445-9.

31.Ram VS, Parthiban, Sudhakar U, Mithradas N, Prabhakar R. Bonebiomarkers in periodontal disease: a review article. J Clin Diagn Res. 2015;9(1):ZE07-10.

32.Prasanthi B, Kannan N, Patil R. Effect of Diuretics on Salivary Flow, Composition and Oral Health Status: A Clinico-biochemical Study. Ann Med Health Sci Res. 2014;4(4):549-53.

33.Leary BK, Leary ME, Bryner S, Pycraft M, Engle R, Olfert IM, et al. Oral Health and Salivary IgA Responses in Division 1 Female Athletes. Int J Exerc Sci Conf Proc. 2022;9(10):49.

34.Gómez SI, Jaramillo LM, Moreno GC, Roa NS, Rodríguez A. Differential reactivity of salivary igA and igG against streptococcus mutans proteins in humans with different caries experience. Acta Odontol Latinoam. 2015;28(1):3-12.

35.Mousavizadeh A, Afroozi B, Hadinia F, Azarshab M, Hadinia A. The Relationship between Salivary and Serum IgA and IgG Levels and Dental Caries in Adults. Clin Lab. 2021;67(8):201209.

36.Needleman I, Ashley P, Petrie A, Fortune F, Turner W, Jones J, et al. Oral health and impact on performance of athletes participating in the London 2012 Olympic Games: a cross-sectional study. Br J Sports Med. 2013;47(16):1054-8.