

The Impact of Fatigue on Physical Statefluctuations in Men Withgenu Varumas Well as Normal Knees

S.K. Mosavi (MSc)*¹, M.R. Eslamipour (MSc)¹, S.S. Shojaeddin (PhD)¹

1.Department of Sports Injury and Corrective Exercises, Faculty of Physical Education and Sport Sciences, Kharazmi University of Tehran, Tehran, I.R.Iran

Received: Jan 2th 2015, Revised: Feb 4th 2015, Accepted: May 6th 2015

ABSTRACT

BACKGROUND AND OBJECTIVE: Genu varum is a risk factor for the knee osteoarthritis onset. Given the role of exhaustion on the physical condition, understanding physical state fluctuations in people with genu varum, will give us a good insight for prevention of knee osteoarthritis. The aim of this study was to compare the effect of exhaustive fatigue on body state fluctuations of men with genu varum and normal knees.

METHODS: This quasi-experimental study was performed on 40 healthy male students, including 20 with genu varum and 20 with normal knees (with the average q angle of 25.5 ± 8.0 and 97.8 ± 86.0 , respectively). Genu varum deformity was measured using a caliper and goniometer. Body fluctuations were recorded using the force measuring diagram, and Strand modified protocol was used to induce fatigue.

FINDINGS: There was no significant difference between the two groups after exhaustion in terms of anterior-posterior stability, but there was a significant decrease in medial-lateral stability after exhaustion (genu varum 86.296 mm/sec and normal knee 38.200 mm/sec, pressure center shifting, respectively); comparing genu varum to the normal knee, these changes were significant ($p=0.04$). Interior-exterior stability in both groups, 10 minutes after exhaustion was fully recovered.

CONCLUSION: Inside displacement of the line of gravity in genu varum sufferers can lead to increased volatility of the lateral posture, and since genu varum deformity causes internal rotation of the leg and turns it into pronation of the subtalar joint in the weight-bearing position, therefore, these changes in foot structures can cause leg function alteration in control of the internal-external (lateral) balance.

KEY WORDS: Fatigue, Posture, Genu varum.

Please cite this article as follows:

Mosavi SK, Eslamipour MR, Shojaeddin SS. The Impact of Fatigue on Physical Statefluctuations in Men Withgenu Varumas Well as Normal Knees. J Babol Univ Med Sci. 2015;17(7):29-35.

* Corresponding Author: S.K. Mosavi (MSc)

Address: Department of Sports Injury and Corrective Exercises, Faculty of Physical Education and Sport Sciences, Kharazmi University, Tehran, I.R. Iran

Tel: +98 21 22258084

Email: kazem_mosavi6486@yahoo.com

Introduction

Daily physical activity and exercise require physical state control, defined as the ability to maintain balance and orientate the body (2, 1); while potential of physical damage increase, is associated with imbalance. Increase in the pressure center swing is considered to be associated with increased prevalence of injuries due to impaired neuro muscular control or balance (2). Sensory posture control input and movement systems are the basic means to maintain physical state (3). Fatigue is also one of the factors influencing body control, that is, if due to local (environmental) exhaustion, any of the visual sensory and atrial systems transfer inaccurate information, or due to general fatigue, central nervous system is disturbed and enforcement is not enough for compensation, the balance would be disrupted (4). Ageberg and colleagues reported that due to loss of the ability to maintain one foot balance, the risk of injury when exhausted is increased (5). Wang and colleagues' research on high school basketball players also showed that most of the lower extremity injuries occur during single leg weight-bearing (6). In a study, Reimer *et al* showed that controlling physical state while in a single-leg standing position, following fatigue, is of great importance (7). Lower limbs are responsible for absorbing pressure during ground contact and adjust the amount of weight (8). Genu varum is among abnormalities in the frontal knee plane with high prevalence among athletes and non-athletes (9). Genu varum shifts the path of the forces from the center of the knee to its internal part and transfers more pressure to the internal structure of the knee, such that the reaction force in the internal sector is around 5.3 times more than the outer part (10). Research shows that on one hand genu varum leads to the loss of cartilage in the tibial joint, and on the other hand leads to the development of osteoarthritis (11). Some studies have mentioned genu varum as a risk factor for patellar femoral pain syndrome and as a predictor of the risk of damage to the ligaments of the knee-joint, including: anterior cruciate ligament injury (anterior cruciate ligament) and posterior cruciate ligament (posterior cruciate ligament) (11, 10). Given that the foot is conjunction of the body and the ground, structural distortions, especially knee injuries have increased risk, and may not be an obstacle to participation in activities by athletes (8). Genu varum abnormalities cause instability in

weight bearing and physical state and changes in the quality of body control (11), by disrupting the orientation of the line of gravity to the base of support (1) and shifting the line of gravity from the center to the inside of the knee (12). Due to the impact of joint structural deformities, especially the knee, in increasing the risk of damage during physical and sport activities, and the need to raise awareness of preventive factors of sport injuries, such as anterior cruciate ligament rupture (6), and increased risk of knee osteoarthritis in patients with genu varum, as well as the limited research performed in the area of the effects of fatigue on the control of physical state in people with genu varum deformity, this study aimed to compare the effect of exhaustive activity fatigue on fluctuations of physical state and balance recovery time in people with genu varum and normal knees.

Methods

This quasi-experimental study was conducted on the male students of Kharazmi University of Tehran, Faculty of Physical Education and Sport Sciences. An initial caliper evaluation of the knee joint was performed on members of the study population. The following factors were regarded as exclusion criteria: professional athlete, injury in the past six months, surgery and injuries in the back and lower extremities, activity limitation on physician's advice, lack of normal strength and full range of motion in the joints of the lower extremities, arthritis or rheumatoid arthritis, impairment due to neuro muscular disorders, actual leg length discrepancy of more than a centimeter through examination of participants, and having other abnormalities such as flat foot, hollow foot, etc. Due to the one-leg stand method of measurement of static balance, the Q angle of the dominant leg in participants was taken into account as eligible. If this angle was less than 8 degrees, it designated knee abnormalities, otherwise if this value was between 8 and 10, it was considered normal (14). Finally, 40 subjects (20 patients with genu varum and 20 patients with normal knees) were selected using non-probability sampling, all of them were matched for age and weight. The feet with which subject kicked the ball was considered the dominant foot. After complete description of the study to the subjects, their consent for participating in research and their personal information such as

age, history of sports, and the number of exercise sessions per week were collected. Using a force plate (model BERTEC, made in USA) physical state fluctuations during the one-leg stand (the dominant) with a sampling frequency of 400 Hz were recorded. In this test, the subjects looked at a certain point. During the test, the hands were placed on the trunk and the knee flexion angle of the non-dominant foot was equal to 90 degrees. Force plate data were recorded for 30 seconds, in order to avoid problems such as insufficient preparation and fatigue, only those incoming data between 5 and 25 seconds were recorded and used for analysis. The Strand altered protocol was used to induce fatigue (5). After an appropriate introduction, each subject was asked to warm up for 3 minutes on a non-inclined treadmill at a speed of 2 miles/hr (3.28 km/h). At the end of the 3 minute warm up, speed was increased to 5 miles/hr (8.2 km/h) for the next 3 minutes. After 3 minutes of running, the slope was increased at a rate of 5.2% every 2 minutes, until the exhaustion protocol ended, in other words up to failure to continue with the protocol (16, 15). To maximize accuracy and minimize fatigue errors, Ageberg pressure perception and heart rate were used.

In accord with this protocol until reaching exhaustion, participant was verbally encouraged to continue running. Minimum heart rate was to reach more than 60% of maximal heart rate and maintained for about 5 min; Ageberg scale was to reach 14–17 (hard or very hard) to halt activities (5). The protocol was chosen because running, creates a situation similar to sports activities and tilt can localize the fatigue in lower limb seven more (5). After suspension of activity, the subjects were required to do a re-tests (first post-test). In order to determine the balance recovery time, every 5 minutes, i.e. 5, 10, 15 and 20 min after the fatigue protocol, static balance test was performed like the pre-test (second, third, fourth and fifth post-tests).

Fluctuations in the body condition were calculated using the Matlab version R2009a

software. To describe the two groups, descriptive statistics, and to analyze the data, inferential statistics, were used. Given the normal distribution of the data, parametric tests were used to analyze the data.

Considering, there were two independent variables in the study group (with 2 significant levels of normal knee and genu varum), and two examination states (with 2 significant levels of pre and post-test) mixed method analysis of variance (Mixed ANOVA) was used. For inter-group comparison, the independent t-test, and for comparing with in groups, paired t-test (comparison tests with the pre-tests) were used, and $p < 0.05$ was considered significant.

Results

In this study, both the genu varum and normal groups had an average age of 75 ± 90.19 and 59.1 ± 20 years, and body mass index was equal to 24.1 ± 37.22 and 42.1 ± 54.22 kg/m², respectively, also the distance between the internal knee breadth was 70.0 ± 65.5 and 99.0 ± 25.1 cm and Q angle was equal to 8.0 ± 25.5 and $86.0 \pm 79.8^\circ$, respectively. Results from comparison of anterior-posterior stability index before and after the test showed that this index was not significant in the genu varum and normal groups.

But the results of the internal-external stability in the pre-test and post-test showed that this index was significant in both genu varum and normal groups ($p = 0.04$) (table 1). Paired comparison of anterior-posterior dynamic stabilization index between pre-test and post-tests showed that there is no significant difference between the two groups (tables 2 and 3). While there was a significant difference in internal-external static stability index between the pre-test and post-test in the normal and genu varum groups. On the other hand no notable difference between the pre-test and the 3rd post-test of the genu varum and three other groups was observed.

Table 1. Before and after comparison of anterior-posterior stability and internal-external static test between the genu varum and normal groups

Groups variable	Mean \pm SD		T	p-value
	genu varum	Normal knee		
Anterior-posterior static stability (mm/s)	26.5 \pm 26.14	37.6 \pm 25.14	83.0	42.0
Internal-external static stability index (mm/s)	45.7 \pm 86.30	83.6 \pm 38.20	18.2	04.*0

Table 2. Before and after comparison of the sagittal and medial-lateral static stability tests in the genu varum group

Variables	The genu varum group					
	Pre-test	Post-test 1	Post-test 2	Post-test 3	Post-test 4	Post-test 5
Anterior-posterior static stability index (mm/s)	52.108	78.113	71.112	51.110	93.109	57.108
within group test	-	23.0	29.0	54.0	67.0	71.0
Internal-external static stability index (mm/s)	44.134	30.164	30.156	77.139	56.136	71.135
Within group test	-	00.*0	00.*0	34.0	49.0	56.0

Table 3. Before and after comparison of the sagittal and medial-lateral static stability test in the normal knee group

Variables	Normal knee group					
	Pre-test	Post-test1	Post-test2	Post-test3	Post-test4	Post-test5
anterior-posterior static stability index (mm/s)	50.113	75.127	27.121	07.115	23.114	63.114
Within group test	-	19.0	23.0	34.0	58.0	79.0
Internal - external static stability index	26.128	64.148	27.142	93.136	78.129	52.128
Within group test	-	00.*0	00.*0	27.0	41.0	68.0

Significance at the 0.05 level

Discussion

This study showed that compared with the pre-test, in the post-test, the anterior-posterior static stability index for the genu varum and normal groups was not significant; also the average differences in the two groups were not significantly different. Nicolas and colleagues proved that lower limb local muscle exhaustion is the main reason for reduction of the balance; they stated the trunk extensor muscle fatigue as the main controlling factor (17). Bearing this in mind, the compensatory mechanisms of the lower limb neuromuscular performance can be the reason behind lack of changes in the balance of the individuals after the fatigue protocol (17).

The Vuillerme et al study showed that the waist neuromuscular system has an important role in controlling the standing posture on the legs, in fact athlete subjects attempt to adapt to fluctuating conditions, and this adaptation occurs in the anterior-posterior level (18). Vuillerme and colleagues, in their research concluded that the stability of the sagittal, in the over fatigue program is not subject to changes, and stated its main reason, in case of general fatigue, as physical state maintenance muscles entering compensatory postural control

before local fatigue mode and holding the balance. They also concluded that in the presence of the visual system the anterior-posterior static variable is affected less (15).

Likewise, Greig et al in their study stated that due to fatigue, anterior-posterior stability, relative to the lateral-index, is less affected, and stated the anatomical shape of bones and soft tissue structure as the cause (19). The results of this study are inconsistent with the results of Reimer et al (20) and Ageberg et al (5), probably due to differences in fatigue protocols and type of induced exhaustion. The present study results showed that the static lateral stability index from the pre-test to the post-test, in both normal and genu varum groups, showed a significant difference and the average change between groups was significant.

It can be concluded that fatigue due to exhaustive activities, decrease lateral-stability in both groups, which was more evident in the genu varum group. Changes in the frontal plane knee alignment, automatically can disrupt the static balance in subjects with normal and symmetrical genu varum deformity, which can alter the joint

weight distribution pattern. This asymmetric distribution of weight in the frontal plane can cause increased posture volatility in knees and ankles (21). Hence, this deformity in leg structure can cause functional changes in the structures set up to control the balance (21). The results of this part of the study are fully compatible with the results of Nardone et al (16) and Letafatkar et al (22) studies.

The present research findings, are partly compatible and partly incompatible with the results of Reimer et al (20). Based on their findings the hip and ankle muscles fatigue (individually), both reduce the state stability in the anterior-posterior and internal-external directions (20). The difference between the two results may be due to the fatigue and exhaustion protocols. Also, Greig and Walker-Johnson, reported lack of significant changes in the stability indices following the fatigue protocol (19). This lack of consistency can be attributed to the differences between research population (semi-professional football players), fatigue protocol (90 minutes of treadmill running) and the test method (single leg standing test on a moving platform).

Ageberg and colleagues also observed increase in range of motion in the sagittal pressure center (anterior-posterior), following cycling, but such a change in the transverse plane (internal-external) did not happen. They attributed the reduced balance to reduced joint receptive activities, muscle spindles and Golgi tendinous organs (5).

According to the present study results, the anterior-posterior static stability state was unchanged, while the stability of the internal-external static condition significantly decreased, and fatigue protocol was recovered within 10 minutes. Nardone et al (16), have reported a significant increase in body fluctuations following exhaustive exercise on the treadmill. One possible reason for this discrepancy is the difference criteria used to stop the fatigue protocol.

It seems that the induced fatigue was more than in the present research and naturally it required more time to recover, it is also possible that enrollment of younger individuals (18–23 years old) in the present study is the reason for faster recovery time. Yaggie and colleagues also showed temporary reduction of state control and a linear relationship between the

direction of oscillation and oxygen consumption in subjects, following physical fatigue (23). Furthermore Bove et al mentioned that fast recovery of the consumed oxygen is one of the reasons for short-term volatility following exhaustion (24). The results of the present research were consistent with Yaggie and colleagues (23) and Bove et al (24), while they were conflicting with Susco and colleagues (25), Khanna et al (4), and Fox and colleagues (26). The reason behind this disparity can be differences in means of balance measurement, fatigue protocols and different post-tests.

Some studies in the field of fatigue have included variables other than balance variables. Baker et al have reported anterior muscle strength recovery within 15 minutes after long-term fatigue protocol, and within 5 minutes after short-term fatigue protocol. They stated that while doing short-term activities, metabolic processes are activated and lead to fast recovery state, while during long term activities it is non-metabolic factors that delay the recovery process (27). Balance recovery following the fatigue before exercise is very important, otherwise the effects of fatigue can lead to a decrease in performance and increase in the risk of injury. Recovery of muscle function depends on severity and duration of exercise and recovery time (4).

It seems that inner shifting line of gravity in people with genu varum can lead to increased lateral posture volatility, thus, this deformity in the leg structures can alter the leg function in balance control. We suggest that this group of people be subject to balance control training exercise programs before physical activity in order to improve balance control. In order to gain a better understanding of the subject matter, it is recommended that a similar study on three groups of genu varum, knock knee and normal be conducted and compared. Because gender affects various factors following fatigue, it is recommended to perform a similar study on women.

Acknowledgments

Hereby we appreciate all Tehran Kharazmi University, Faculty of Physical Education and Sport Sciences laboratory staff and students participating in the study.

References

- 1.Hrysomallis C, McLaughlin P, Goodman C. Balance and injury in elite Australian footballers. *Int J Sports Med*. 2007;28(10):844-7.
- 2.McGuine TA, Greene JJ, Best T, Leverson G. Balance as a predictor of ankle injuries in high school basketball players. *Clin J Sport Med*. 2000;10(4):239-44.
- 3.Reiman BL, Myers JB, Lephart SM. Sensorimotor system measurement techniques. *J Athl Train*. 2002;37(1):85-98.
- 4.Pallavi Kh, Kapoor G, Kalpana Z. Balance deficits and recovery timeline after different fatigue protocols. *Indian j phy occup*. 2008;2(3):42-54.
- 5.Ageberg E, Robert D, Holmstorm E, Friden T. Balance in single-limb stance in healthy subjects – reliability of testing procedure and the effect short duration sub-maximal cycling. *BMC Musculoskelet Disord*. 2003; 4(14):1-16.
- 6.Wang HK, Chen CH, Shiang TY, Jan MH, Lin KH. Risk-factor analysis of high school basketball-player ankle injuries: a prospective control cohort study evaluating postural sway, ankle strength, and flexibility. *Arch Phys Med Rehabil*. 2006;87(6):821-5.
- 7.Reimer RC, Wikstrom EA. Functional fatigue of the hip and ankle musculature cause similar alterations in single leg stance postural control. *J Sci Med Sport*. 2010;13(1):161-6.
- 8.Mc Clay IS, Hamill J, Buchanan TS. Lower extremity kinematics and kinetics differences in runners with high and low arches. *J Appl Biomech*. 2001;17(2):153-63.
- 9.Hadadnezhad M, Letafatkar A. The relationship between genu varum abnormality and lower extremity's performance and strength in teenage footballers. *J Rehabil Sci*. 2011;7(2):188-96. [In Persian]
- 10.Sharma L, Song J, Dunlop D, Felson D, Lewis CE, Segal N, et al. Varus and valgus alignment and incident and progressive knee osteoarthritis. *Ann Rheum Dis*. 2010;69(11):1940-5.
- 11.Van Gheluwe B, Kirby KA, Hagman F. Effects of simulated genu valgum and genu varum on ground reaction forces and subtalar joint function during gait. *J Am Podiatric Med Assoc*. 2005;95(6):531-41.
- 12.Levangie PK, Norkin CC. Joint structure & function: A Comprehensive Analysis, 5th ed. F.A Davis Company; 2011.p.1-588.
- 13.Samaei A, Bakhtiary AH, Elham F, Rezasoltani A. Effects of genu varum deformity on postural stability. *Int J Sports Med*. 2012; 33(6):469-73.
- 14.Smith TO, Hunt NJ, Donell ST. The reliability and validity of the Q-angle: a systematic review. *Knee Surg Sports Traumatol Arthrosc*. 2008;16(12):1068-79.
- 15.Vuillerme N, Burdet C, Isableu B, Demetz S. The magnitude of the effect of calf muscles fatigue on postural control during bipedal quiet standing with vision depends on the eye-visual target distance. *Gait Posture*. 2006;24(2):166-72.
- 16.Nardone A, Tarantola J, Giorano A, Schieppatti M. Fatigue effects on body balance. *Electroencephalogr Clin Neurophysiol*. 1997; 105(4):309-20.
- 17.Vuillerme N, Anziani B, Rougier P. Trunk extensor muscle fatigue affects undisturbed postural control in young healthy adults. *Clin Biomech (Bristol, Avon)*. 2007;22(5):489-94.
- 18.Vuillerme N, Pinsault N, Chenu O, Fleury A, Payan Y, Demongeot J. Postural destabilization induced by trunk extensor muscle fatigue is suppressed by use of a plantar pressure based electro-tactile biofeedback. *Eur J Appl Physiol*. 2008;104(1):119-25.
- 19.Greig M, Walker-Johnson C. The influence of soccer-specific fatigue on functional stability. *Phys Ther Sport*. 2007;8(4):185-90.

- 20.Reimer R, Wikstrom E. Functional fatigue of the hip and ankle musculature cause similar alterations in single leg stance postural control. *J Sci Med Sport*. 2009;13(1):161-6.
- 21.Bakhtiary AH, Fatemi E, Rezasoltani A. Genu varum deformity may increase postural sway and falling risk. *Koomesh*. 2012;13(3): 330-8.
- 22.Letafatkar K, Alizadeh MH, Kordi MR. The effect of exhausting exercise induced fatigue on the double-leg balance of elite male athletes. *J Soc Sci*. 2009;5(4):445-51.
- 23.Yaggie J, Armstrong WJ. Effect of lower extremity fatigue on indices of balance. *J Sport Rehabil*. 2004;13(4):312-22.
- 24.Bove M, Faelli E, Tacchino A, Lofrano F, Cogo CE, Ruggeri P. Postural control after a strenuous treadmill exercise. *Neurosci Lett*. 2007;418(3):276-81 .
- 25.Susco T, Valovich MLT, Gansneder B, Shults S. Balance recovers within 20 minutes after exertion as measured by the balance error scoring system. *J Athl Train*. 2004;39(3):241-6 .
- 26.Fox CR, Paige GD. Effect of head orientation on human postural stability following unilateral vestibular. *J Vestib Res*. 1990-91;1(2): 153-60.
- 27.Baker AJ, Kostov KG, Miller RG, Weiner MW. Slow force recovery after long duration exercise: metabolic and activation factors in muscle fatigue. *J Appl Physiol*(1985). 1993;74(5):2294-300.