# The Relationship between Physical Activity during Pregnancy and Postpartum Mood in Primiparous Women

# M. Mirghafourvand (PhD)<sup>1</sup>, S. Mohammad-Alizadeh Charandabi (PhD)<sup>2</sup>, M. Asghari Jafarabadi (PhD)<sup>3</sup>, A. Mohammadi (MSc)<sup>4</sup>, S. Soltanpour Gharibdoosti (MSc)<sup>\*5</sup>

Department of Midwifery, Faculty of Nursing and Midwifery, Tabriz University of Medical Sciences, Tabriz, I.R.Iran
 Social Determinants of Health Research Center, Faculty of Nursing and Midwifery, Tabriz University of Medical Sciences, Tabriz, I.R.Iran
 Road Traffic Injury Research Center, Faculty of Health, Tabriz University of Medical Sciences, Tabriz, I.R.Iran
 Department of Midwifery, Faculty of Nursing and Midwifery, Arak University of Medical Sciences, Arak, I.R.Iran
 Students' Research Committee, Faculty of Nursing and Midwifery, Tabriz University of Medical Sciences, Tabriz, I.R.Iran

## J BabolUniv Med Sci; 18(6); Jun 2016; PP:35-41 Received: Jan 6<sup>th</sup> 2016, Revised: May 1<sup>th</sup> 2016, Accepted: May 9<sup>th</sup> 2016

#### ABSTRACT

**BACKGROUND AND OBJECTIVE:** Physical activity might reduce postpartum depressive symptoms and improve temperament. This study aimed to evaluate the relationship between physical activity during pregnancy and postpartum mood in primiparous women.

**METHODS:** This cohort study was conducted in 165 primiparous women aged 18-35 years referring to the healthcare centers in Tehran, Iran, during 2013-2014. The participants were chosen through stratified random sampling and divided into two groups of low physical activity (n=68) and moderate or high physical activity (n=97). Both groups completed the International Physical Activity Questionnaire (IPAQ) during weeks 28 and 34 and Depression Anxiety Stress Scale (DASS) at the end of the sixth postpartum week. For each sub-scale, the minimum and maximum possible scores of this scale are 0 and 21, respectively.

**FINDINGS:** Mean total scores of stress, anxiety, and depression of the low physical activity group were  $9.85\pm5.74$ ,  $5.61\pm5.11$ , and  $6.23\pm5.77$ , respectively, while for the moderate or high physical activity group they were  $9.88\pm5.84$ ,  $5.72\pm5.03$ , and  $6.51\pm5.70$ , respectively. In addition, no significant difference was observed between the two groups in terms of mean total score of stress (p=0.969), anxiety (p=0.585), and depression (p=0.396) at the end of the sixth postpartum week. Moreover, no statistically significant relationship was observed between the level of physical activity during pregnancy and postpartum mood.

**CONCLUSION:** According to our results, physical activity during pregnancy is not associated with postpartum stress, anxiety, and depression.

**KEY WORDS:** Physical activity, Stress, Anxiety, Depression, Pregnancy, Postpartum.

#### Please cite this article as follows:

Mirghafourvand M, Mohammad-Alizadeh Charandabi S, Asghari Jafarabadi M, Mohammadi A, Soltanpour S. The Relationship between Physical Activity during Pregnancy and Postpartum Mood in Primiparous Women. J Babol Univ Med Sci. 2016;18(6):35-41.

Address: Students' Research Committee, Faculty of Nursing and Midwifery, Tabriz University of Medical Sciences, Tabriz, I.R.Iran Tel: +98 41 34796770 E-mail: sedigheh.soltanpour@yahoo.com

## Introduction

**P**hysical activity is defined as any body movement performed by skeletal muscles and consuming sufficient energy, which should not be misconstrued as sport. Physical activity incorporates activities such as playing, walking, doing housework, gardening, and dancing (1). On the other hand, physical inactivity is considered as the fourth leading risk factor for global mortality (2). Temperament is a stable feeling, generally affecting the perception of individuals about the world and their function. In general, variations in postpartum mood are categorized as grief, depression, and postpartum psychosis based on the severity, onset time, and duration of symptoms (3). While mental health problems in the first trimester have been evaluated in the majority of high-income countries, few studies have been conducted on this issue in lowand middle-income countries (4). In developing countries, major mental health issues have been observed in one in of 3-5 women during pregnancy and postpartum periods; however, this rate was estimated at one in 10 women in developed countries (5). According to the literature, stress during pregnancy can be associated with some degrees of mental illness (6). Mental health issues, including depression and anxiety, are considered as the most common mental health disorders (5, 7) and are some of the most current problems during and after pregnancy worldwide (7).

Evidence suggests that 10-15% of women are diagnosed with mental disorders, especially anxiety that is one of the most common mental problems (8). Vulnerability to depression increases during puerperium (9). It seems that primiparous women might be at an increased risk of fatigue, distress, anxiety, frustration, and depression in early postpartum period (10). Since these symptoms affect 6.5-12.9% of women in the first postpartum year (12), it seems necessary to conduct relevant interventions to address family issues, as well as social and personal functions in women suffering from depressive symptoms (13).

According to the previous studies, postpartum depression is one of the most serious health risks and a common disorder adversely affecting mothers, neonates, and families (11). Therefore, promoting awareness regarding the relationship between physical activity and anxiety might help physicians recognize those women at risk of postpartum fatigue, distress, or even depression (14).

Part of our understanding regarding the effects of physical activity on postpartum mood is based on some

previous studies conducted on the impact of physical activity on postpartum depressive symptoms (15-18). Norman et al. conducted a study to evaluate the effects of therapeutic exercise program and healthcare education on maternal mental health. However, the results were indicative of a significant improvement in depression scores in the intervention group (19). In a study by MacDonald, conducted on 130 Canadian primiparous women, no significant association was observed between the level of physical activity during pregnancy and temperament at the sixth and 12<sup>th</sup> postpartum weeks (3).

activity could improve Enhanced physical postpartum maternal health (20, 21). Symons Downs et al. concluded that physical activity could significantly decrease postpartum depressive symptoms, enhance mood, and increase energy and endurance during and after pregnancy (22). Therefore, participation in physical activities to fight against common postpartum mental and physical disorders seems to be of paramount importance (23). Given the short- and longterm benefits of physical activity for overall health, it seems necessary to conduct studies on the impact of physical activity on temperament during and after pregnancy (24), Based on the review of available literature, no studies were found on the relationship between physical activity during pregnancy and its effects on postpartum mood in Iran. Since maternal, fetal, and family health is highly affected by behavioral symptoms, this study was conducted to investigate the relationship between physical activity during pregnancy and postpartum mood.

#### **Methods**

This cohort study was conducted on 165 primiparous pregnant women after obtaining approval of Ethics Committee. The samples were divided into two groups of low physical activity and moderate or high physical activity. In this study, the inclusion criteria were 1) aged 18-35 years, 2) gestational age of 28-34 weeks, 3) Iranian nationality, 4) willingness to participate in the study, 5) having a phone, 6) singleton pregnancy, and 7) residence in Tehran. Meanwhile, the only exclusion criterion was being diagnosed with diseases majorly limiting physical activity (24, 26).

Based on the study by MacDonald et al. (3) on the relationship between physical activity and stress, anxiety, and depression, sample size was estimated at 66 with odds ratio of 3.2, 95% confidence interval, and

80% power using the G-Power software. However, the final sample size was considered 165 regarding cluster sampling design (double) effect and predicting 25% attrition (27). In the pilot study, approximately 14% of the samples had low physical activity. Given the aim of the present study and the importance of balancing the samples to provide the required test power, sampling was continued until selection of 55 subjects with low physical activity. Sampling was performed in healthcare centers affiliated to Shahid Beheshti University of Medical Sciences in Tehran, Iran, during 2013-2014. With regard to the six districts of Tehran, 12 centers and 10 healthcare stations were selected using stratified random sampling. Afterwards, the samples were selected from the list of pregnant women with gestational age of 28-34 weeks using the same sampling method.

After contacting the subjects via phone calls and explaining the objectives of the study, they were asked to visit the healthcare centers to complete the questionnaires in case of willingness to participate and meeting the inclusion criteria. Written informed consent was obtained from the participants at the first visit, and the questionnaires were completed by the subjects thereafter. If the predetermined sample size was not achieved in the two groups, the same process pursued. After six weeks of postpartum follow-up, the samples visited in person and completed the DASS by interview.

A researcher-made demographic questionnaire, IPAQ, and DASS were employed to collect data. The short version of IPAQ includes seven items, completed by the participants during pregnancy and at the sixth postpartum week.

Data were collected in terms of type of physical activity performed by primiparous women (as part of their daily routines) in the past seven days (28). First, metabolic equivalents were calculated for each physical activity at 3.3, 4, and 8 for walking, moderate, and intense physical activities, respectively. Then, these numbers were multiplied by the number of days and duration of the mentioned physical activity (in minute) (29). Validity and reliability of DASS were confirmed by Baghbani Moghadam et al. at the Cronbach's alpha of 0.7, which is indicative of a good internal consistency. Reliability of this questionnaire was also established using Spearman-Brown prediction formula (0.9) (30). Stress, depression, and anxiety of the samples were evaluated using the selfreporting DASS-21 scale at the sixth postpartum week.

This scale consists of 21 items and three sub-scales including stress, depression, and anxiety (seven items in each sub-scale), rated using a 5-point Likert scale ranging from never (zero) to a lot () (31). The minimum and maximum scores of each sub-scale were zero and 21, respectively (32)This scale has also been validated in Iran and its reliability has been confirmed, as well (33). Content and face validities were used to approve the validity of the study tools. In addition, its reliability was demonstrated in terms of internal consistency (Cronbach's alpha coefficient) and replicability (intraclass correlation [ICC]) using testretest method and 30 subjects. Cronbach's alpha coefficient and ICC (95% confidence interval) were estimated at 0.93 and (0.96-0.81) 0.90, respectively. Data analysis was performed in SPSS version 21 and normality of quantitative variables was evaluated using skewness and kurtosis.

The results were indicative of normal distribution of stress, anxiety, and depression scores. Mann-Whitney U and Chi-squared tests were run to investigate the relationship between physical activity and total scores of stress, depression, and anxiety. In addition, P-value of less than 0.05 was considered statistically significant.

#### Results

The stages of the present study are presented in figure 1.



Figure 1. Study procedure

According to the results, samples of the two groups were homogenous in terms of demographic characteristics (table 1). Mean total score of physical activity during pregnancy was  $2563.7\pm3414.6$  (table 2). Mean values for stress, anxiety, and depression during pregnancy were  $9.8\pm5.6$ ,  $5.5\pm5.3$ , and  $5.8\pm5.9$ , respectively, while in the moderate and high physical activity group, they were  $9.88\pm5.84$ ,  $5.72\pm5.03$ , and  $6.51\pm5.70$ , respectively. Nevertheless, no significant difference was observed between the groups regarding the total mean scores of stress (p=0.969), anxiety (p=0.585), and depression (p=0.396) at the end of the sixth postpartum week.

According to our findings, no statistically significant link was observed between the level of physical activity during pregnancy and stress (p=0.823), anxiety (p=0.820), and depression (p=0.807) (table 3).

# Table 1. Evaluation of the relationship between demographics of pregnant women with low and moderate or high physical activity referring to the healthcare centers in Tehran, Iran at the sixth postpartum week

Physical activity	Low	Moderate or high	<b>P-value</b>
	( <b>n=68</b> )	( <b>n=97</b> )	
Body Mass Index (kg/m <sup>2</sup> )			
≤19.8	7(10.3)	8(8.2)	*0.501
19.8-25.9	40(58.8)	62(63.9)	
26.0-29.9	12(17.6)	21(21.6)	
≤30.0	9(13.2)	6(6.2)	
Occupational status			
Housewife	63(93)	87(90)	#0.516
Employed	5(7.4)	10(10.3)	
Educational status			
Elementary	4(6)	6 (6.2)	*0.784
Junior and senior high school	10(15)	21(22)	
Diploma	39(57.4)	39(40.2)	
Academic education	15(22.1)	31(32)	
Sufficient income for living expenses			
Sufficient	5(7.4)	23(24)	*0.032
To some extent	53(78)	62(64)	
Insufficient	10(15)	12(12.4)	
Number of family members			
Two	54(79.4)	83(86)	#0.300
More than two	14(21)	14(21)	
Housing status			
Personal residence	17(25)	29(30)	<sup>#</sup> 0.717
Rental or family	51(75)	68(70.1)	
*** Preferable feeding mode			
Breastfeeding only	61(90)	79(81.4)	#0.306
Breast milk with bottle	7(10)	18(18.5)	

\*Chi-squared test process, \*Chi-squared test

\*\*\*\*There was only one case of bottle-feeding in the group of moderate or high physical activity.

 Table 2. Total score of physical activity in women with low and moderate or high physical activity referring to

 the healthcare centers in Tehran. Iran

the neutrical c centers in Tenran, fran					
Physical activity	*Mean±SD	Median	25-75%	Domain	
Total score	2536.72±3413.6	1188	436-3360	0-28476	
Low	1680.93±2204.73	775	267.50-1731	0-9786	
Moderate or high	3182.58±3946.54	1680	706.50-5113	0-28476	

			D
Physical activity	Low	Moderate or high	<b>P-value</b>
	(N=63)	(N=84)	
Stress			
Low (0-7)	22(35)	32 (37)	*0.823
Moderate(8-14)	26(41)	34(39.5)	
Severe(≥15)	15(24)	20(23.5)	
Mean±SD	$9.8 \pm 5.6$	9.9±5.8	#0.969
Anxiety			
Low(0-7)	45(72.0)	60(70.0)	*0.820
Moderate(8-14)	14(22.0)	20(23.0)	
Severe(≥15)	4(6.0)	6(7.0)	
Mean±SD	$5.5 \pm 5.3$	5.7±5.0	#0.585
Depression			
Low (0-7)	42(66.0)	53(62.0)	*0.807
Moderate(8-14)	13(21.0)	24(28.0)	
Severe(≥15)	8(13.0)	9(10.0)	
Mean+SD	$5.8 \pm 5.9$	6.5+5.7	#0.396

Table 3. The relationship between mood at sixth postpartum week and low and moderate or high physical activity in women referring to healthcare centers in Tehran. Iran

\* Chi-squared test process, \* Mann-Whitney U test

#### **Discussion**

According to the results of the present study, no significant relationship was observed between physical activity and postpartum stress, anxiety, and depression. The current study is the first cohort study aiming to appraise the association between physical activity during pregnancy and postpartum mood in Tehran, Iran. Armstrong demonstrated attenuated depressive symptoms and improved adjustment in the intervention group with walking program (17), which was not in line with outcomes of the present study. The discrepancy between our results and those obtained by Armstrong and another study by Radloff could be due to homogenizing participants in terms of depression level before the evaluation of physical activity level and its association with postpartum depression, whereas we determined the level of physical activity during pregnancy and evaluated postpartum mood (34). Although previous reports have exhibited the positive effect of physical activity on postpartum depression (3), temperament, energy, endurance (22), and common mental and physical disorders during pregnancy (29), in the current study, no significant relationship was found between physical activity during pregnancy and postpartum mood. This inconsistency in results might be due to the different times of mood appraisal, which was performed in the sixth postpartum week in this study, and the 12<sup>th</sup> postpartum week in other studies. It seems that this 12week interval is suitable for the samples to adjust to the new event in their life (i.e., being a mother) and might attenuate stress, anxiety, and even depression. Another study by Downs et al. conducted on 74 mothers in postpartum period was indicative of improved mood due to physical activity. This discrepancy in results might be due to the difference in time of mood evaluation. In the current study, this evaluation was conducted on the sixth postpartum week, while it was performed in the fifth postpartum month in the aforementioned study. It seems that this 12-week interval was a suitable for the samples to adjust to their new event in life (i.e., being a mother) and might lead to their decreased levels of stress and anxiety (22). The main limitations of our study were short time interval (26-28th week of gestation to sixth postpartum week) and lack of adequate follow-up during the mentioned period. Therefore, future studies are recommended to be conducted over longer periods of time with prolonged follow-ups and shorter time intervals. Moreover, this study was carried out on women residing in Tehran, which might not be generalized to residents of rural areas of this city or even other cities in Iran and women with underlying diseases. Given the importance of physical activity for promotion of overall health, conducting clinical trials is recommended to investigate the effect of regular physical activity during pregnancy on postpartum mood to gain an accurate understanding of the relationship between physical activity during pregnancy and postpartum mood.

#### Acknowledgements

Hereby, we wish to thank the Deputy of Research and Technology and School of Nursing and Midwifery of Tabriz University of Medical Sciences, Tabriz, Iran, and chancellors of Tehran, Shahid Beheshti, and Iran universities. We would also like to thank the participants for their cooperation.

#### References

1.WHO. Global strategy on diet, physical activity and health. 2004. Available from: http://www.who.int/dietphysicalactivity/en/. 2.WHO. Global status report on noncommunicable diseases 2010. Available from: http://www.who.int/nmh/publications/ncd\_report\_full\_en.pdf.

3.Macdonald IB. Physical activity and postpartum functional status in primiparous women. Kingston: Queen's University; 2011. Available from https://qspace.library.queensu.ca/bitstream/1974/6812/1/Barbacsy-MacDonald\_Ibo\_201109\_MSc.pdf

4.WHO. Maternal mental health and child health and development in low and middle countries. 2008 Available from http://www.who.int/mental\_health/prevention/suicide/mmh\_jan08\_meeting\_report.pdf.

5.WHO. Maternal mental health and child health and devolopment in resource-constrained setting. 2007 ;[Available from: http://apps.who.int/iris/bitstream/10665/70118/1/WHO\_RHR\_09.24\_eng.pdf

6.NasiriAmiri F, Salmalian H, Hajiahmadi M, Ahmadi A. Association between prenatal anxiety and spontaneous preterm birth. J Babol Univ Med Sci. 2009;11(4):42-8.[In Persian]

7.Alipour Z, Lamyian M, Hajizadeh E. Anxiety and fear of child birth as predictors of postnatal depressionin nulliparous women. Women Birth. 2011;25(3):37-43.

8.NasiriAmiri F, Salmalian H, Hajiahmadi M. Association between prenatal anxiety and spontaneous preterm birth. J Babol Univ Med Sci. 2009;11(4):42-8.[In Persian].

9. Stowe Z, Hostetter AL, Newport DJ. The onset of postpartum depression: Implications for clinical screening in obstetrical and primary care. Am J Obstet Gynecol. 2004;192(2):522-6.

10.Morse C. Postnatal depression, issues in midwifery practice today. The 2<sup>nd</sup> national midwifery forum. Australia: LaTrobe University; 1991.p.72-6.

11.Hosseini H, Naghibi AH, Khademloo M. Post partum depression and its relationship with some related factors. J Babol Univ Med Sci. 2008; 10(2): 76-81. [In Persian]

12.Gaynes BN, Gavin N, Meltzer-Brody S, Lohr KN, Swinson T, Gartlehner G. Perinatal depression: prevalence, screening accuracy, and screening outcomes.Evid Rep Technol Assess (Summ). 2005;119:1-8.

13.Posmontier B. Functional status outcomes in mothers with and without postpartum depression. J Midwifery Womens Health. 2008;53(4):310-18.

14.McVeigh CA. Anxiety and functional status after child berth. Aust Coll Midwives Inc J. 2000;13(1):14-8.

15.Dritsa M, Dupuis G, Khalifé S. Effects of a home-based exercise intervention on fatigue in postpartum depressed women: results of a randomized controlled trial. Ann Behav Med. 2008;35(2):179-87.

16.Heh SS, Huang LH, Ho SM, Fu YY, Wang LL. Effectiveness of an exercise support program in reducing the severity of postnatal depression in taiwanese women. Birth. 2008;35(1): 60-5.

17. Armstrong K, Edwards H. The effects of exercise and social support on mothers reporting depressive symptoms: A pilot randomized controlled trial. Int J Ment Health Nurs. 2003;12(2): 130-8.

18. Armstrong K, Edwards H. The effectiveness of a pram-walking exercise programme in reducing depressive symptomatology for postnatal women. Int J Nurs Pract. 2004;10(4):177-94.

19.Norman E, Sherburn M, Osborne R, Galea M. An exercise and education program improves well-being of new mothers: A randomized controlled trial. Phys Ther. 2010;90(3): 348-55.

20.Baheiraei A, Mirghafourvand M, Mohammadi E, Mohammad-AlizadehCharandabi S. The experiences of women of reproductive age regarding health-promoting behaviours: A qualitative study. BMC Public Health. 2012;12:573-83.

21.Baheiraei A, Mirghafourvand M, Mohammadi E, Mohammad-AlizadehCharandabi S, Nedjat S. Determining appropriate strategies for improving women's health promoting behaviours: using the nominal group technique. East Mediterr Health J. 2013;19(5):409-16.

22.Symons Downs D, Hausenblas HA. Women's exercise beliefs and behaviors during their pregnancy and postpartum. J Midwifery Womens Health. 2004;49(2):138-44.

23.Gjerdingen DK, Froberg DG, Chaloner KM, McGovern PM. Changes in women's physical health during the first postpartum year. Arch Fam Med. 1993;2(3):277-83.

24.Davies G, Wolfe L, Mottola M, MacKinnon C. Joint SOGC/CSEP clinical practice guideline: Exercise in pregnancy and the postpartum period. Can J Appl Physiol. 2003;28(3):330-41.

25.Artal R, O"Toole M. Guidelines of the American college of obstetricians and gynecologists for exercise during pregnancy and the postpartum period. Br J Sports Med. 2003;37(1):6-12.

26.Warburton D, Nicol C, Bredin S. Health benefits of physical activity: The evidence. CMAJ. 2006;174(6):801-9.

27.Salganik MJ. Variance estimation, design effects, and sample size calculations for respondent-driven sampling. Urban Health. 2006;83(7):98-112.

28. Craig C, Marshall A, Sjostrom M, Bauman A, Booth M, Ainsworth B. International physical activity questionnaire: 12-country reliability and validity. Med Sci Sports Exerc. 2003;35(8):1381-96.

29. Booth ML, Ainsworth BE, Pratt M, Ekelund ULF, Yngve A, Sallis JF, et al. International physical activity questionnaire: 12-country reliability and validity. Med Sci Sports Exerc. 2003;195:1381-95.

30.BaghianiMoghaddam MH, BakhtariAghdam F, AsghariJafarabadi M, Allahverdipour H, DabaghNikookheslat S, Safarpour Sh. The Iranian version of international physical activity questionnaire (IPAQ) in Iran: content and construct validity, factor structure, internal consistency and stability. World Appl Sci. 2012;18(8):1073-80.

31. Bastard J, Tiran D. Aromatherapy and massage for antenatal anxiety: Its effect on the fetus. Complement Ther Clin Pract. 2009;15(4):230-3.

32. Henry JD, Crawford JR. The short-form version of the depression anxiety stress scales (DASS: 21): construct validity and normative data in a large nonclinical sample. Br J Health Psychol. 2005;44(2):227-39.

33.Sahebi A, Asghari M, Salari R. Validation of depression anxiety and stress scale (DASS-21) for an Iranian population. Iran Psychol. 2005;4:299-313. [In Persian]

34.Radloff, L, Sawyer L. The CES-D scale: A selfreport depression scale for research in the general population. Appl Psychol Meas. 1977:1(3) 385-401.