

Analgesic Activity of the Extract of Aerial Parts of Colza (*Brassica Napus*) in Mice

S. Kazemi (PhD)¹, A. Shirvani (MD)¹, M. Hashemi (MSc)¹, A.A. Moghadamnia (PhD)^{*1,2}

1.Department of Pharmacology, Babol University of Medical Sciences, Babol, I.R.Iran

2.Neuroscience Research Center, Babol University of Medical Sciences, Babol, I.R.Iran

J Babol Univ Med Sci; 18(5); May 2016; PP: 38-43

Received: Feb 24th 2016, Revised: Mar 2th 2016, Accepted: Mar 15th 2016.

ABSTRACT

BACKGROUND AND OBJECTIVE: It has been reported that Cruciferous family plants such as *Brassica napus* (Colza) are abundant sources of flavonoid compounds that involve in prostaglandin synthesis and may show analgesic and anti-inflammatory effects. The purpose of this study was to investigate the analgesic effect of hydroalcoholic extract of aerial parts of Colza in comparison with morphine, with or without naloxone in mice.

METHODS: The male mice weighing 18-26 g were divided into experimental groups (6 mice in each group) and received i.p. injections of 50, 100, 250 and 500 mg/kg of the hydroalcoholic extract of colza and morphine with or without naloxone, respectively. Normal saline was used as control. The hot-plate test was performed to evaluate the analgesic effects of all treatments and pain latency was measured at 15, 30, 45, 60, 75, and 90 minutes after injection of the drugs.

FINDINGS: Pain tolerance of the mice receiving various doses of colza extract was significantly increased compared to the control. Moreover, simultaneous injection of morphine (20 mg/kg) and colza extract (250 mg/kg) increased pain tolerance compared to morphine alone. Also, simultaneous injection of colza extract and morphine at 10 minutes after naloxone (1 mg/kg), increased analgesia in the animals. The highest analgesia was observed after treatment with morphine and colza extract (250 mg/kg) at 30 minutes after the injection (32 ± 2 seconds).

CONCLUSION: According to the results, the hydroalcoholic extract of the aerial parts of colza induced analgesia in mice during the hot-plate test. This effect may be attributed to the presence of flavonoid compounds in the extract, which confirms the analgesic properties of colza.

KEY WORDS: Colza (*Brassica napus*), Flavonoids, Morphine, Naloxone, Inflammation, Analgesic.

Please cite this article as follows:

Kazemi S, Shirvani A, Hashemi M, Moghadamnia AA. Analgesic Activity of the Extract of Aerial Parts of Colza (*Brassica Napus*) in Mice. J Babol Univ Med Sci. 2016;18(5):38-43.

*Corresponding author: A. Moghadamnia (PhD)

Address: Center Research for Neuroscience, Babol University of Medical Sciences, Babol, I.R.Iran

Tel: +98 11 32199596

E-mail: moghadamnia@yahoo.com

Introduction

Pain as an unpleasant feeling, is the reaction of the body to deleterious stimulants and works as a protective mechanism. Many medications have been introduced to fulfill the goal of pain relief but the adverse drug reactions can limit their administration. It is believed that complications are rare when natural plants are used instead of chemical medications, therefore, they are in the center of scientists' attention as future analgesics.

The plants of Brassica family including *brassica Napus* are known for their anticancer activities (1-9). They have various chemical compounds such as polyphenols that have anti-inflammatory properties. It has been shown that *brassica oleracea* can be used in treating gout, jaundice, rheumatoid arthritis, obesity, and kidney diseases (10). Analgesic and anti-inflammatory effects of *brassica oleracea* were previously reported (11). It was reported *brassica juncea* can reduce gastric pain induced by acetic acid (12). Some other characteristics of this family have been studied by investigators but analgesic properties of *brassica napus* (Colza in Farsi) have not been studied before. Colza contains thiol, indol, and flavonoid compounds (11, 12).

It also contains phenolic compounds which can reduce inflammation and oleic and linoleic acids which regulate steroid's metabolism (7, 13). This plant is broadly cultivated in the north of Iran. The oil of colza seeds has been used in food industries. This study was done to evaluate the analgesic effects of the different doses of hydroalcoholic extract of Colza aerial parts in mice

Methods

The research protocol was approved by the ethics committee of Babol University of Medical Sciences and no animal was hurt during this study. Male albino mice weighing 18 to 26 g were used. The mice were divided into experimental groups and different doses of colza, morphine, or naloxone were intraperitoneally injected (i. p.). Control groups were only injected with normal saline. At least, six mice were used in each treatment or control groups. To prepare Colza extract, the plants were collected in spring from the farms in northern areas of Iran. The aerial parts were washed, dried, and then chopped and powdered. Ethyl alcohol

80% was then added to this powder and filtered after 72 hours. The filtered liquid was left for 72 hours for evaporation the ethanol and Colza extract to remain. Colza has been administered in four different doses (50, 100, 250 and 500 mg/kg) and the dose of 250 mg/kg of colza extract was selected to use with morphing and naloxone.

Naloxone (1mg/kg) as opioid antagonist was injected 10 min before morphine or doses of colza. Hot-plate latency times were measured for each animal before any injection and 15, 30, 45, 60, 75 and 90 minutes after drug or saline treatments. Hot plate device was set to 54 °C and end point was defined as the moment when the animal started to lick his legs. Cut-off point was set at 40 seconds. If each animal stayed on hot plate without any reactions for more than 40 seconds, the mouse was excluded from the experiment. ANOVA post-post hoc Tukey test was used to compare the data. P-value ≤ 0.05 was considered significant.

Results

Table 1 summarizes data of the treatments groups and presents results of pain latency time in hot-plate test as mean and standard deviation. The pain latency time of colza (50, 100, 250, and 500 mg/kg) groups and saline groups is presented in figure 1. Group receiving colza 500 mg/kg shows a significant pain latency time compared to saline at 15 minute after injection ($p=0.004$). Evaluation of the concomitant use of morphine and colza (250 mg/kg) showed that injection with morphine 20 mg/kg can significantly increase pain latency at times 15 ($p=0.001$), 60 (0.01), and 90 ($p=0.018$) minutes after injection in comparison with the injection of the same concentration of morphine alone (fig 2). It seems that combination of colza and morphine has better analgesic effects than single morphine injection. In figure 2, the slope of the curve of the effect of Colza (250 mg/kg) + morphine (20 mg/kg) versus time is ascending after 90 minutes while the morphine (20 mg/kg) alone is descending at the same time. Naloxone (1 mg/kg) was used as a pretreatment of colza (250 mg/kg) and morphine (20 mg/kg) the results showed significant differences at 15($p<0.001$), 30($p=0.006$), 60($p<0.001$), and 90 ($p=0.002$) minutes after injection compared to group receiving colza 250 mg/kg or morphine 20mg/kg(fig 3).

Table 1. Mean (SD) of hot-plate pain latency time (sec) in mice receiving doses of colza in comparison to saline and morphine. Pre-treatment naloxone (1 mg/kg) was used as opioid antagonist (the number of animals in each group were at least 6 mice)

Treatment groups	Time	Baseline	15 min	30 min	45 min	60 min	75 min	90 min
Saline		10.52(5.56)	13.04(8.11)	14.52(5.51)	12.20(8.19)	10.56(6.16)	11.16(5.56)	11.40(7.66)
Colza 250 mg/kg		12.08(9.11)	14.00(7.48)	11.25(4.92)	13.83(7.72)	12.00(6.67)	12.50(6.74)	9.50(2.78)
Colza 500 mg/kg		10.93(4.98)	18.00(9.20)	13.56(6.28)	12.20(5.90)	11.53(7.33)	13.40(6.40)	9.83(4.03)
Morphine (20 mg/kg)		15.22(8.66)	23.72(8.66)	22.38(9.97)	20.50(11.65)	16.39(8.85)	17.67(10.20)	15.72(5.65)
Colza 250 + Morphine (20 mg/kg)		14.08(6.92)	30.92(8.70)	23.58(11.73)	20.83(11.14)	25.42(11.28)	19.25(9.72)	21.33(7.89)
Naloxone+Colza250+Morphine20		17.67(12.19)	29.00(11.21)	32.33(10.52)	17.83(11.58)	22.33(14.45)	23.00(13.94)	27.33(14.08)
Naloxone+Colza500+Morphine20		16.67(2.42)	28.50(9.16)	28.50(24.83)	24.83(9.22)	23.67(8.21)	18.50(7.69)	19.67(4.08)

*Pretreatment with naloxone (1 mg/kg) used as morphine antagonist (minimum of six animals in each group); **Animals receiving colza extract (500 mg/kg) had a significant difference with normal saline group at 15 minutes after injection ($p=0.004$);***No significant differences observed in other groups




Figure 1. Mean of pain threshold in hot-plate test (sec) in mice receiving colza extract at different doses (50, 100, 250 and 500 mg/kg) compared to control group.

*Animals administered with colza extract (500 mg/kg) had a significant difference with control group at 15 minutes after injection ($p<0.001$).




Figure 2. Mean of pain threshold in hot-plate test (sec) in mice receiving colza extract (250 mg/kg) and morphine (20 mg/kg) at different times (simultaneous and separate administration).*Simultaneous administration of morphine (20 mg/kg) and colza extract (250 mg/kg) had a significant difference with morphine only ($p<0.001$).




Figure 3. Mean of pain threshold in hot-plate test (sec) at different times in mice receiving colza extract (250 mg/kg) with morphine (20 mg/kg) and naloxone (1 mg/kg) (10 minutes before injection) and mice receiving naloxone and morphine (30 mg/kg)

*Animals administered with naloxone, colza extract (250 mg/kg) and morphine (20 mg/kg) had a significant difference with mice receiving colza extract (250 mg/kg) and morphine ($P < 0.001$); **Significant differences between mice receiving colza extract and morphine with normal saline group.

Discussion

Medicinal plants have been used for relieving pain for a long time (11, 14). In this study, analgesic effects of hydroalcoholic extract of aerial parts of colza were investigated using hot-plate test in mice. This study confirmed that colza, in comparison to saline, shows analgesic activity in high dose at 15 minutes after i.p. injection. On the other hand, colza can increase analgesic activity of morphine in almost all times of the measurements. Naloxone as a specific opioid antidote could not fully antagonize the morphine effect when colza was added. It seems that the mechanism of action of colza may be different and opioid involvement in its action cannot be confirmed (11). In addition, it has been found that colza can maintain analgesia for more than 90 minutes. Antioxidants have various functions in the body (15, 16), and flavonoid compounds have been shown to have remarkable anti-inflammatory and analgesic properties (17, 18). These compounds are abundantly found in the leaves and roots of the species of the Brassica family (19). Flavonoids exert their analgesic effects by crossing the blood-brain barrier and influencing the α -2 adrenergic and GABA receptors. Moreover, these compounds inhibit the enzymatic activity of cyclooxygenase-2 in damaged tissues preventing the production of prostaglandins (20, 21). Several studies have suggested that through inhibiting the NMDA receptors, flavonoids reduce calcium concentration in the cells, which diminishes the enzymatic activity of nitric oxide synthase and phospholipase A2. This mechanism might explain the analgesic properties of flavonoids through decreasing the activity of nitric

oxide and prostaglandins (22). Previous studies have confirmed that flavonoids are able to decrease metabolism of arachidonic acid and production of prostaglandin E through degradation of tumor necrosis factor (TNF- α) and inhibition of cyclooxygenase (23). Furthermore, some flavonoids, such as apigenin, prevent the imbalance of lipid signaling pathways (24, 25). Previous studies have shown the analgesic properties of some species of the Brassica family, such as *Brassica juncea*, which may exert anti-inflammatory effects through the decrease of prostaglandin synthesis (26, 27). However, there is lack of adequate data regarding the exact analgesic effects of colza. According to the literature, the most important compounds in *Brassica oleracea* are phenols and flavonoids, which are known to have anti-inflammatory and analgesic properties (28, 29). In conclusion, findings of the present study indicated that the effect of *Brassica napus* on the analgesic properties of morphine was due to the presence of flavonoids in this medicinal plant. Based on the results, the hydroalcoholic extract of *Brassica napus* could be an effective complementary medicine for pain relief. Nevertheless, it is recommended that further investigations be conducted in this regard in order to clarify the exact mechanism of action of colza.

Acknowledgments

This investigation was financially supported by the Deputy of Research and Technology of Babol University of Medical Sciences.

References

1. Michaud DS, Spiegelman D, Clinton SK, Rimm EB, Willett WC, Giovannucci EL. Fruit and vegetable intake and incidence of bladder cancer in a male prospective cohort. *J Natl Cancer Inst.* 1999; 91(7):605-13.
2. Spitz MR, Duphorne CM, M.A. Detry, Pillow PC, Amos CI, Lei L, Andrade M et al. Dietary intake of isothiocyanates: evidence of a joint effect with glutathione S-transferase polymorphisms in lung cancer risk. *Cancer Epidemiol Biomarkers Prev.* 2000;9(10):1017-20.
3. Lewis S, Brennan P, Nyberg F, Ahrens W, Constantinescu V, Mukeria A, et al., Cruciferous vegetable intake, GSTM1 genotype and lung cancer risk in a non-smoking population. *Iarc Sci Publ.* 2002;156:507-8.
4. Zhang, SM, Hunter DJ, Rosner BA, Giovannucci EL, Colditz GA, Speizer FE et al., Intakes of fruits, vegetables, and related nutrients and the risk of non-Hodgkin's lymphoma among women. *Cancer Epidemiol Biomarkers Prev.* 2000; 9(5):477-85.
5. Cohen JH, Kristal AR, Stanford JL. Fruit and vegetable intakes and prostate cancer risk. *J Natl Cancer Inst.* 2000;92(1):61-8.
6. Kolonel LN, Hankin JH, Whittemore AS, Wu AH, Gallagher RP, Wilkens LR et al. Vegetables, fruits, legumes and prostate cancer: a multiethnic case-control study. *Cancer Epidemiol Biomarkers Prev.* 2000; 9(8):795-804.
7. Terry P, Wolk A, Persson I, Magnusson C. Brassica vegetables and breast cancer risk. *JAMA.* 2001; 285(23):2975-7.
8. Du XZ, Ge XH, Yao XC, Zhao ZG, Li ZY Production and cytogenetic characterization of intertribal somatic hybrids between *Brassica napus* and *Isatis indigotica* and backcross progenies. *Plant Cell Rep.* 2009; 28(7):1105-13.
9. Farag MA, Sharaf Eldin MG, Kassem H, Abou el, Fetouh M. Metabolome classification of *Brassica napus* L. organs via UPLC-QTOF-PDA-MS and their anti-oxidant potential. *Phytochem Anal.* 2013; 24(3):277-87.
10. Zaidi SF, Aziz M, Muhammad JS, Kadowaki M. Review: Diverse pharmacological properties of *Cinnamomum cassia*: A review. *Pak J Pharm Sci.* 2015;28(4):1433-8.
11. Carvalho CA, Fernandes KM, Matta SL, Silva MB; Licursi de Oliveira L, César Fonseca C. Evaluation of antiulcerogenic activity of aqueous extract of *Brassica oleracea* var. capitata (cabbage) on Wistar rat gastric ulceration. *Arq Gastroenterol.* 2011;48(4):276-82.
12. Wu GX, Lin YX, Ou MR, Tan DF. [An experimental study(II) on the inhibition of prostatic hyperplasia by extract of seeds of *Brassica alba*]. *Zhongguo Zhong Yao Za Zhi.* 2003, 28(7):643-6.
13. Tzagoloff A. Metabolism of Sinapine in Mustard Plants. I. Degradation of Sinapine into Sinapic Acid & Choline. *Plant Physiol.* 1963; 38(2):202-6.
14. Radha Krishnan K, Babuskin S, Azhagu Saravana Babu P, Sasikala M, Sabina K, Archana G, Sivarajan M, et al. Antimicrobial and antioxidant effects of spice extracts on the shelf life extension of raw chicken meat. *Int J Food Microbiol.* 2014;171:32-40.
15. Shahaboddin ME, Pouramir M, Moghadamnia AA, Parsian H, Lakzaei M, Mir H. *Pyrus biossieriana* Buhse leaf extract: An antioxidant, antihyperglycaemic and antihyperlipidemic agent. *Food Chem.* 2011;126(4):1730-3.
16. Mahjoub S, Tamaddoni A, Zanjanchi-Nikoo M, Moghadamnia AA. The effects of beta-carotene and vitamin E on erythrocytes lipid peroxidation in beta-thalassemia patients. *J Res Med Sci.* 2007;12(6):301-7. [In Persian]
17. Fatouros NE, Pashalidou FG, Aponte Cordero WV, van Loon JJ, Mumm R, Dicke M, et al, Anti-aphrodisiac compounds of male butterflies increase the risk of egg parasitoid attack by inducing plant synomone production. *J Chem Ecol.* 2009;35(11):1373-81.

18. Babaee N, Moslemi D, Khalilpour M, Vejdani F, Moghadamnia Y, Bijani A, et al. Antioxidant capacity of calendula officinalis flowers extract and prevention of radiation induced oropharyngeal mucositis in patients with head and neck cancers: a randomized controlled clinical study. *Daru*. 201;21(1):18.
19. Cartea, M.E., M. Francisco, P. Soengas, Velasco P. Phenolic compounds in brassica vegetables. *Molecules*. 2011;16(1):251-80.
20. Hitz WD, Carlson TJ, Booth JR Jr, Kinney AJ, Stecca KL, Yadav NS. Cloning of a higher-plant plastid omega-6 fatty acid desaturase cDNA and its expression in a cyanobacterium. *Plant Physiol*. 1994;105(2):635-41.
21. Morteza-Semnani K, Saeedi M, Hamidian M, Vafamehr H, Dehpour AR. Anti-inflammatory, analgesic activity and acute toxicity of *Glaucium grandiflorum* extract. *J Ethnopharmacol*. 2002;80(2-3):181-6.
22. Rotelli AE, Guardia T, Juárez AO, de la Rocha NE, Pelzer LE. Comparative study of flavonoids in experimental models of inflammation. *Pharmacol Res*. 2003;48(6):601-6.
23. dos Santos MD, Almeida MC, Lopes NP, de Souza GE. Evaluation of the anti-inflammatory, analgesic and antipyretic activities of the natural polyphenol chlorogenic acid. *Biol Pharm Bull*. 2006. 29(11):2236-40.
24. Lopes LS, Pereira SS, Silva LL, Figueiredo KA, Moura BA, Almeida FR, Sousa FC. Antinociceptive effect of topiramate in models of acute pain and diabetic neuropathy in rodents. *Life Sci*. 2009;84(3-4):105-10.
25. Okada Y, Okada M. Protective effects of plant seed extracts against amyloid beta-induced neurotoxicity in cultured hippocampal neurons. *J Pharm Bioallied Sci*. 2013;5(2):141-7.
26. Lapa Fda R, Gadotti VM, Missau FC, Pizzolatti MG, Marques MC, Dafré AL, Farina M, Rodrigues AL, Santos AR. Antinociceptive properties of the hydroalcoholic extract and the flavonoid rutin obtained from *Polygala paniculata* L. in mice. *Basic Clin Pharmacol Toxicol*. 2009;104(4):306-15.
27. dos Santos DA, Fukui Mde J, Dhammika Nanayakkara NP, Khan SI, Sousa JP, Bastos JK, Andrade SF, et al. Anti-inflammatory and antinociceptive effects of *Baccharis dracunculifolia* DC (Asteraceae) in different experimental models. *J Ethnopharmacol*. 2010; 127(2):543-50.
28. Leal LK, Ferreira AA, Bezerra GA, Matos FJ, Viana GS. Antinociceptive, anti-inflammatory and bronchodilator activities of Brazilian medicinal plants containing coumarin: a comparative study. *J Ethnopharmacol*. 2000;70(2):151-9.
29. An S, Han JI, Kim MJ, Park JS, Han JM, Baek NI, et al. Ethanolic extracts of *Brassica campestris* spp. rapa roots prevent high-fat diet-induced obesity via beta(3)-adrenergic regulation of white adipocyte lipolytic activity. *J Med Food*. 2010;13(2):406-14.