

The Effect of Kangaroo Mother Care on Mental Health of Mothers with Premature Infants

Y. Zahedpasha (MD)¹, A. Salarmanesh (MSc)², S. Khafri (PhD)³, S. Mouodi (MD)⁴, A. Arzani (PhD)*⁵

1.Non-Communicable Pediatric Disease Research Center, Health Research Institute, Babol University of Medical Sciences, Babol, I.R.Iran

2.Student Research Committee, Babol University of Medical Sciences, Nursing &Medwifery Faculty, Babol, I.R.Iran

3.Infertility and Reproductive Health Research Center, Health Research Institute, Babol University of Medical Sciences, Babol, I.R.Iran

4.Neuroscience Research Center, Health Research Institute, Department of Psychiatry, Babol University of Medical Science, Babol, I.R.Iran

5.Nursing Care Research Center, Health Research Institute, Babol University of Medical Sciences, Babol, I.R.Iran

J Babol Univ Med Sci; 20(6); June 2018; PP: 7-13

Received: Dec 26th 2017, Revised: Mar 5th 2018, Accepted: Apr 10th 2018.

ABSTRACT

BACKGROUND AND OBJECTIVE: Mothers with premature infants experience high levels of psychological stress. Considering the importance of maternal mental health as a parent-child care provider, this study was conducted to evaluate the effectiveness of Kangaroo Mother's Care (KMC) of premature infant on mother's mental health.

METHODS: In this quasi-experimental study, 60 mothers with premature infants (less than 37 weeks pregnant) admitted to Ayatollah Rohani Hospital, Babol University of Medical Sciences, were selected by non-random sampling method. They divided into two study groups marked as experimental group (n=30) and control group (n=30). Data were collected using the demographic questionnaire and General Health of Goldberg & Hiller questionnaire. The 28-item questionnaire consisted of 4 sub-scales: physical symptoms, anxiety and sleep symptoms, social function, and depression symptoms. A score of over 22 indicates a symptom of the disease. In the intervention group, mothers spent at least 2 hours, 3 to 4 times a day for one week. The control group had the current process in the care. The questionnaire was completed at the beginning and one month after the admission of the newborn in both groups and the data were compared.

FINDINGS: Finding showed that the overall mental health score in the intervention group before and after maternal care was (28.77±13.09) to (13.66±6.5) and in the control group (30.86±15.97) to (20.99±8.89) that this difference was significant (p<0.01). In relation to the scales in the intervention and control groups, respectively, the anxiety and sleep patterns (3.7±1.5 in contrast to 5.67±3.39), social function disorder (5.9±2.58 vs 7.21±2.2) Physical symptoms (2.13±1.88 vs. 5.2±3.31) had a significant difference between the two groups (p<0.05), but in the case of depressive symptoms (1.03±0.3) vs 2.7±2.31), this difference was not significant (p>0.05).

CONCLUSION: Regarding the results of this research, use of Kangaroo Mother's Care (KMC) for improve of premature infant on mother's mental health is step for health family.

KEY WORDS: *Kangaroo Mother's Care (KMC), Mental Health, Premature.*

Please cite this article as follows:

Zahedpasha Y, Salarmanesh A, Khafri S, Mouodi S, Arzani A. The Effect of Kangaroo Mother Care on Mental Health of Mothers with Premature Infants. J Babol Univ Med Sci. 2018; 20(6): 7-13.

* Corresponding Author: A. Arzani (PhD)

Address: Faculty of Nursing &Medwifery, Babol University of Medical Sciences, Babol, I.R.Iran

Tel: +98 11 32190597

E-mail: Arzaniaf@gmail.com

Introduction

During pregnancy and childbirth, tremendous changes have taken place in women's spirits, which make them very sensitive to psychological stimuli and sometimes cause problems (1). The process of coping with maternal delivery is naturally challenging (2). Premature infants (less than 37 weeks of gestation) constitute 11.4% of the births of the newborns (3). The premature birth of an infant affects the physical and emotional-psychological well-being of parents (4) and presents them with many problems, and can negatively affect the mood of their mothers and their reciprocal patterns of behavior with their infants. (5). Among the factors influencing the mental health of mothers after delivery is the health status of the born infant. (6). The results of the researches show that 40-76% of mothers with preterm and low birth weight infants have an experience of depression and other psychological symptoms including incompatibility (8,7). The need to care of a premature infant puts mothers in an unpredictable and unexpected condition (9). The hospitalization experience of these infants in the intensive care unit is scary for mothers (10). The longer period and the duration of care for the infants in the Neonatal Intensive Care Unit (NICU), results in the greater anxiety and stress for mothers, and the mental health of mothers will be further affected (11). Usually, an incubator or warmer device is used to care for these infants. While these devices often unnecessarily remove infants from their mothers and deprive them of skin-to-skin contact, hence, Kangaroo Mother Care (KMC), in which the infant is in skin contact with mother, is introduced as an appropriate alternative to incubation care of premature infants who overcome the initial problems. So today, this type of care is recommended in accordance with the recommendations of the WHO and the Neonatal Health Department, which is to be implemented continuously. The experience of implementing such a project, despite the provision of a KMC package has been seriously implemented in a limited number of university centers, and is still not fully institutionalized at the national level (12). Reports demonstrated that in the KMC method, mental stress, stress and doom of mothers' conscience are less than the conventional one, and mothers prefer to use skin contact. They have expressed more self-confidence, higher self-esteem and a sense of satisfaction and ability

to do positive work for their premature infants (13, 14). Findings of the study by Faramarzi et al showed that maternal care using KMC method had increased maternal compatibility with existing problems. In other words, this method of care will help mothers to have a better performance in infant care in stressful conditions due to the birth of premature and low birth weight infants. In another study, De Macedo and colleagues reported that KMC plays an important role in mood changes and reduction of depression in mothers (15). Kangaroo Mother Care (KMC) also improves the health and well-being of infants through more effective control of body temperature, success in breastfeeding, infection prevention, and closer relationship between mother and child (16), and prepare them to establish a to-way and coordinated interactive pattern (17). A review of the literature showed that most research focused on the effect of Kangaroo Mother Care on the physical condition of the infant (19,18), and in the few studies, mothers with premature infants were as target population. Since the underlying psychological basis of the individual comes from the very early childhood and infancy, primary support of the mother is very important in future interactions between them (20), and on the other hand, given the broad and valuable view of prevention of mental health problems, in this study, the effect of Kangaroo Mother Care on the mental health of mothers with premature infants was investigated.

Methods

This quasi-experimental study was carried out using available random sampling method after obtaining permission from the Ethics Committee of Babol University of Medical Sciences with number: Mubabol.REC.1394.119. This study was performed on 60 mothers with premature infant admitted to the Neonatal Intensive Care Unit at the Medical and Educational Center of Ayatollah Rouhani affiliated to Babol University of Medical Sciences in 2012-2013 in two groups of control and intervention. Mothers with premature infants (less than 37 weeks of gestational age) admitted to the neonatal intensive care unit who had a tendency to perform Kangaroo Mother Care and their infants had a stable physiological status and weight of between 1000 and 2500 grams, at least reading and writing skills in Persian, singleton, pregnancy with

tendency, no history of depression and birth of premature infants, no abnormalities in the infant and no adverse events were included in the study. Discharge earlier than one week after starting the Kangaroo Mother Care was exclusion criteria of the study. In this research, with 95% confidence, the test power of 80% and $d = 0.65$ (21), the sample size was estimated to be 30 in each group. After cooperation of mothers to participate in the study and following all ethical considerations in the research (confidentiality of information, obtaining written consent from mothers, voluntary participation in the study) demographic information questionnaire and Goldberg & Hiller questionnaire with 28 questions were used as a tool for data collection (20). The mothers were trained in the intervention group to put the infant naked in the middle of their chest to make skin contact, and provide maternity care for at least 2 hours, 3-4 times a day for one week. Monitoring the mother's performance was continuously and effectively performed by the researcher to ensure that the intervention was carried out. However, the mothers in the control group were in the process of ongoing care. The mental health of mothers first and after one month after admission was considered based on the questionnaire. The General Health Questionnaire, a 28-item questionnaire, was developed by Goldberg and Hiller in 1979, which includes four levels of physical symptoms, anxiety and sleep disorders, impaired social function, and severe depression (20).

Of the 28 items of the questionnaire, items 1 to 7 are related to the scale of physical symptoms, 8 to 14 symptoms of anxiety and sleep disorders, and 15 to 21 items are related to the assessment of social function symptoms and ultimately 22 to 28 items measure the symptoms of depression. The questionnaire, as a screening tool, spends around 10-12 minutes, (22) examines the person's mental status in a recent month (23). In this study, 28 questions were used in the standard form and responses were coded using Likert score (0-1-2-3). The cut-off point in this study was that individuals with a score of 22 and below were enrolled in the healthy group and those with a score of 23 and higher were categorized in the group of people with a disorder (25, 24). Noorbala et al., in evaluating the validity of the general health questionnaire with 28-items, based on traditional scoring, reported cut off score of 6 and based on simple Likert score, reported the

cutoff point of 22, and the reliability coefficient of the questionnaire was 91% (26). In this study, the reliability coefficient was 89%. The questionnaire was completed two times, before the starting of Kangaroo Mother Care at the beginning of the infant admission and after one month of infant admission. This study was performed by comparing the questionnaires completed before and after the intervention by the mother. After collecting the final data, obtained data were analyzed using SPSS 14 software and multivariate covariance analysis (MANCOVA), $p < 0.05$ was considered significant.

Results

In this study, 60 mothers were enrolled. Most of the participating mothers had a high school diploma (65.7%) and 76.6% were housekeeper (Table 1).

Table 1. Distribution of absolute and relative frequency of some demographic characteristics of mothers in two groups

group	variable	intervention N(%)	control N(%)	X ²
Education	Under diploma	2(3.4)	8(13.4)	4.49
	Diploma	17(28.4)	13(21.3)	
	Bachelor	5(8.3)	2(3.4)	
	Higher than diploma	6(10)	7(11.8)	
Occupation	House keeper	25(41.6)	21(35)	2.79
	Employee	5(8.4)	9(15)	

The mean of the most subscales of the intervention group after the implementation of the maternal care program was reduced, so that the overall mental health score in the intervention group before and after maternal care was 28.77 ± 13.09 and 13.66 ± 6.51 , respectively and in the control group was changed from 30.86 ± 15.97 to 20.99 ± 8.89 ($p < 0.01$) (Table 2). The intervention group compared with the control group after receiving maternal care was significantly reduced in the sub variables of the mental health scale, except for the symptoms of depression in the intervention group. In other words, the reduction of mean in variables of the mental health scale for the intervention group by maternal care was statistically significant. The results of the multivariate covariance analysis test for subscales of mental health test after removing the effect of pre-test by hotelling test showed that in the sub variables of the

mental health scale, there was a significant decrease in the mean of depression scores in the intervention group (Table 3). Two groups were similar and did not have a

significant difference in terms of interventional variables such as infant's age, infant's disease during admission, birth rate and age of mothers ($p < 0.05$).

Table 2. Comparison of mental health variables in two groups of intervention and control

Variable	Intervention group			Control group		
	Pre-test Mean±SD	Post-test Mean±SD	p-value	Pre-test Mean±SD	Post -test Mean±SD	p-value
Symptoms of anxiety and sleep disorders	4.62±8.23	1.50±3.70	<0.001	5.31±8.83	3.39±5.67	<0.001
Social function	3.31±9.32	2.58±5.90	<0.001	2.38±10.16	2.20±7.13	<0.001
Symptoms of depression	1.90±3.87	1.30±1.03	<0.001	2.79±3.77	2.31±2.70	<0.002
Physical symptoms	4.27±7.73	1.58±2.93	<0.001	4.85±8.10	3.31±5.20	<0.002
mental health	13.09±28.77	6.51±13.56	<0.001	15.97±30.86	8.89±20.90	<0.001

* Significant difference in the level of 0.05 between the two groups

Table 3. Summary of single-variable covariance test results

Variable	SS	df	MS	F	Sig
Symptoms of anxiety and sleep disorders	47.87	1	47.87	6.99	0.01
Social function	18.78	1	18.78	3.86	0.04
Symptoms of depression	4.88	1	4.88	1.23	0.17
Physical symptoms	57.35	1	57.35	7.17	0.01
mental health	442.52	1	442.52	8.15	0.01

Discussion

The results of this study showed that Kangaroo Mother Care had a significant effect on improving the mental health of mothers in both the whole scale and some of its subscales. There was a significant difference between the mean scores of the subjects in the intervention and control groups at the post-test stage in the whole scales of mental health and its subscales, including physical symptoms, insomnia and anxiety and social function. In confirmation of our findings, the study by Badiie et al. also showed that Kangaroo Mother Care had a positive effect on the mental health of mothers with low-birthweight infant (11). Palizvan et al. also found that kangaroo mother care was effective on the general health of mothers with a neonate (term) and without history of hospitalization (27). Along with the results of our study in the anxiety subscale, Keshavarz et al. showed that skin contact is effective on the positional anxiety of mothers admitted to the cesarean section (28). However, in our study, more intervention days and effect of KMC for a longer period (one month

after admission of infant) have been investigated and evaluated. The study of Karimi et al. showed the effect of mother and infant skin contact on the attachment and anxiety of mothers regarding the health status of the newborn (17). Faramarzi et al. in their study acknowledged that close proximity to the skin and the contact of the skin to the mother and the baby caused a greater sense of comfort, health and compatibility in mothers (2). In the study of Feldman et al., mothers showed more positive emotions with skin contact with their infants and became more compatible with their infant. According to their findings, the rate of depression in mothers was also reduced. In three months of infancy, parents showed more sensitivity to caring for the child and provided a better home environment for them. and in the six-month follow up of mental development, these children developed significantly on the basis of the Bayley development test of infants and the psycho-motor development index (29). Nyqvist et al., in a study in Sweden regarding the effect of Kangaroo Mother Care, found that this intervention reduced the

symptoms of depression in mothers and made parents more sensitive to the baby. In addition, it increases the breastfeeding duration of mothers and leads to better interaction between parents (30).

In a study by Saatsaz et al., the daily contact between mother and the infant for 20-30 minute days for 10 days reduced the severity of postpartum mourning in mothers (31). De Alecar and colleagues in their study emphasized that Kangaroo Mother Care would have an effect on the improvement of postpartum depression, feeling of strength, happiness, and speed of action in the mother (32). In another study, De Macedo and colleagues reported that Kangaroo Mother Care plays an important role in mood change and reduction of depression in mothers (15). In our study, the subscales of depression symptoms did not show a meaningful relationship between the two groups, although the mean scores of this area decreased after intervention. As mentioned, effective components in mental health are symptoms of depression and people with depression symptoms have low self-confidence and self-esteem (33). Therefore, interventions that lead to improving the level of depression in mothers are considered important. In this regard, Lee et al. suggested that Kangaroo Mother Care, in addition to fixing the physiological symptoms of the baby, it also increases the self-esteem of the mothers (13).

However, Arzani et al showed that there was no significant difference in maternal self-esteem score before and after KMC interventions and at the time of discharge (34), which could be related to the type of instrument used to measure the level of self-esteem, the duration of the KMC in a day or the time of completing the self-esteem questionnaire (during the discharge of the baby, which is an anxious and stressful time for parents). Considering the birth rate of the premature infant in the country and the complications of the illness and the admission of the infant to the mother, it is recommended to provide a suitable and non-prescriptive approach that is easy to do and requires no equipment

and high cost. Therefore, it is desirable to provide this care method for mothers with premature infants in the very early days of the infant's birth, and the health care staff and nurses to carry out and promote this care continuously (24 hours), because the mothers are responsible for care and maintenance of the infants and this is something that nurses must understand and prioritize. The limitations of this study are the cultural and social characteristics of the community, which requires people to be conservative in many areas such as mental health, especially in our research, which is used by self-report scales to measure this structure. Therefore, it is suggested that in future research, other appropriate tools be used to measure more accurately the mental health of individuals.

Our research showed that Kangaroo Mother Care can provide more mental health for mothers, leading to reduced physical symptoms, insomnia and anxiety, social ineffectiveness and depression in mothers with preterm infants. Therefore, it is recommended that the result of this research be of interest to maternal and neonatal health care providers in order to continually use this care method in neonatal intensive care units because the mother is in direct interaction with the newborn and, of course, the quality of this relationship and interaction is also effective on the process of infant growth. In addition, the findings of this study can help nursing managers to improve the knowledge of nurses through the provision and holding of workshops, panel discussions and discussions on Kangaroo Mother Care.

Acknowledgment

Hereby, we would like to thank to the Vice-Chancellor for Research and Technology of Babol University of Medical Sciences for financial support and from the colleagues of the neonatal intensive care unit of the Ayatollah Rohani Hospital in Babol which helped us in the process of implementation of this research.

References

1. Korja R, Savonlahti E, Haataja L, Lapinleimu H, Manninen H, Piha J, et al. Attachment representations in mothers of preterm infants. *Infant Behav Dev.* 2009;32(3):305-11.
2. Faramarzi S, Badiie Z, Miri T, Khodadad Z. The effectiveness of kangaroo mother's care (KMC) method of low birth weight infants on maternal adjustment. *Iran J Nurse Res.* 2014;8(4):29-41. [In Persian].
3. Evreklian M, Posmontier B. The impact of kangaroo care on premature infant weight gain. *J Pediat.* 2017;34:10-6.
4. Flacking R, Lehtonen L, Thomson G, Axelin A, Ahlqvist S, Moran VH, et al. Closeness and separation in neonatal intensive care. *Acta Paediat.* 2012;101(10):1032-7.
5. Tooten A, Hall RA, Hoffenkamp HN, Braeken J, Vingerhoets AJ, Van Bakel HJ. Maternal and paternal infant representations: a comparison between parents of term and preterm infants. *Infant Behav Dev.* 2014;37(3):366-79.
6. Trombini E, Surcinelli P, Piccioni A, Alessandrini R, Faldella G. Environmental factors associated with stress in mothers of preterm newborns. *Acta paediat.* 2008;97(7):894-8.
7. Brett JI, Staniszevska S, Newburn M, Jones N, Taylor L. A systematic mapping review of effective interventions for communicating with, supporting and providing information to parents of preterm infants. *BMJ Open.* 2011;1(1):e000023.
8. Jotzo M, Poets CF. Helping parents cope with the trauma of premature birth: an evaluation of a trauma-preventive psychological intervention. *Pediat.* 2005;115(4):915-9.
9. Lindberg B, Ohrling K. Experiences of having a prematurely born infant from the perspective of mothers in northern Sweden. *Int J Circumpolar Health.* 2008;67(5):461-71.
10. Leonard A, Mayers P. Parents' lived experience of providing kangaroo care to their preterm infants. *Health SA Gesondheid.* 2008;13:16-28.
11. Badiie Z, Faramarzi S, MiriZadeh T. The effect of kangaroo mother care on mental health of mothers with low birth weight infants. *Adv Biomed Res.* 2014;3:214.
12. Vakilian R, Haidarzadeh M. Kangaroo Mother Care(KMC service package). Tehran: Ideapardazan Fan & Art Co. 2013. [In Persian].
13. Lee J, Bang KS. The effects of kangaroo care on maternal self-esteem and premature infants' physiological stability. *Korean J Women Health Nurs.* 2011;17(5):454-62.
14. Ahn HY, Lee J, Shin HJ. Kangaroo care on premature infant growth and maternal attachment and post-partum depression in South Korea. *J Trop Pediatr.* 2010;56(5):342-4.
15. de Macedo EC, Cruvinel F, Lukasova K, D'Antino ME. The mood variation in mothers of preterm infants in kangaroo mother care and conventional incubator care. *J Trop Pediatr.* 2007;53:344-6.
16. Jagadale S, Salunkhe J. To evaluate the effectiveness of kangaroo mother care on low birth weight babies. *Int J Sci Res.* 2014;3(8):2017-9.
17. Karimi A, Fatemeh T, Khadivzadeh T, Aghamohammadian Sharbaf HR. The effect of skin to skin contact immediately after delivery on the maternal attachment and anxiety regarding infant. *Iran J Obs, Gynecol Infertil.* 2013;16(67):7-15.
18. Valizadeh S, Hosseini M B, Asghari Jafarabadi M, Mohebbi L. The effect of kangaroo mother care on feeding tolerance in preterm infants. *J Babol Univ Med Sci.* 2014;16(12):16-22. [In Persian].
19. Bera A, Ghosh J, Singh AK, Hazra A, Mukherjee S, Mukherjee R. Effect of kangaroo mother care on growth and development of low birth weight babies up to 12 months of age: a controlled clinical trial. *Acta Paediatrica.* 2014;103(6):643-50.
20. Börjesson B1, Paperin C, Lindell M. Maternal support during the first year of infancy. *J Adv Nurs.* 2004;45(6):588-94.
21. Karimi F Z, Khadivzadeh T, Saeidi M, Bagheri S. The effect of kangaroo mother care immediately after delivery on mother-infant attachment 3 months after delivery. *Int J Pediatr.* 2016;4(9):3561-70.
22. Goldberg DP, Gater R, Sartorius N, Ustun TB, Piccinelli M, Gureje O, et al. The validity of two versions of the GHQ in the WHO study of mental illness in general health care. *Psycholog Med.* 1997;27(01):191-7.
23. Goldberg DP, Hillier VF. A scaled version of the General Health Questionnaire. *Psychological Med.* 1979;9(01):139-45.
24. Ashtekar SV, Kulkarni MB, Sadavarte VS, Ashtekar RS. Analysis of birth weights of a rural hospital. *Ind J Com Med.* 2010;35(2):252.

25. Balaji K, Sankar S, Nandagopal B. Low birth weight of newborns: Magnitude of the problem seen in a 100 bed hospital of a rural area in Vellore district, Tamil Nadu (India). *Ind J Com Med*. 2010;35(2):362.
26. Noorbala A, Mohammad K. The validation of general health questionnaire-28 as a psychiatric screening tool. *Hakim Res J*. 2009;11(4):47-53. [In Persian].
27. Palizvan N, Hekmat K, Sharifi N, JavadiFar N, Haghighizadeh M H, Mousavi P. Effect of kangaroo mother care on postpartum maternal general health status. *Iran J Obstet, Gynecol Infertil*. 2014;17(121):10-17. [In Persian]
28. Keshavarz M, Norozi F, Sayyed Fatemi N, Haghani H. Effect of skin to skin contact on maternal state anxiety in a cesarean section ward. *Knowledge Health*. 2011;5(4):1-6. [In Persian].
29. Feldman R, Eidelman AI, Sirota L, Weller A. Comparison of skin-to-skin (kangaroo) and traditional care: parenting outcomes and preterm infant development. *Pediat*. 2002;110(1):16-26.
30. Nyqvist KH, Anderson GC, Bergman N, Cattaneo A, Charpak N, Davanzo R, et al. Towards universal kangaroo mother care: recommendations and report from the first european conference and seventh international workshop on kangaroo mother care. *Acta Paediat*. 2010;99:820-6.
31. Saatsaz S, Rezaei R, Sharifnia H, Kheirkhah F, Moulookzadeh S, Haji Hosseini F. Effect of mother and newborn skin to skin contact on postpartum blues. *J Babol Univ Med Sci*. 2011;13(3):59-65. [In Persian]
32. de Alencar AEMA, Arraes LC, de Albuquerque EC, Alves JGB. Effect of kangaroo mother care on postpartum depression. *J Tropical Pediat*. 2009;55(1):36-8.
33. Benjamin J, Virginia A, Pedro R. *Psychology books kaplan&sadoks*. 10th ed. Wolters Kluwer Health; 2017. Vol 1, pp: 1606
34. Arzani A, Zahedpasha Y, Ahmadpour KM, Khafri S, Khairkhah F, Aziznejad P. Kangaroo care effect on self-esteem in the mothers of low birth weight infants. *J Babol Univ Med Sci*. 2012;14(3):52-8.